

PESTICIDE ATLAS

Facts and figures about toxic chemicals in agriculture

2022

NIGERIA EDITION



HEINRICH
BÖLL
STIFTUNG



Friends of
the Earth
Europe



BUND
FRIENDS OF THE EARTH GERMANY



Pesticide
Action
Network
Europe

IMPRINT

The **PESTICIDE ATLAS 2022 – NIGERIA EDITION** is jointly published by
Heinrich-Böll-Stiftung, Abuja, Nigeria
Heinrich-Böll-Stiftung, Berlin, Germany
Friends of the Earth Europe, Brussels, Belgium
Bund für Umwelt und Naturschutz, Berlin, Germany
PAN Europe, Brussels, Belgium

Chief executive editors:
Lisa Tostado, Heinrich-Böll-Stiftung European Union (project management)
Dr. Silke Bollmohr, EcoTrac Consulting
Jochen Luckscheiter and Ikenna-Donald Ofoegbu (Nigeria Edition)

Managing editor, graphics research: Martin Eimermacher
Art direction, graphic development: STOCKMAR+WALTER Kommunikationsdesign
Graphic development (Nigeria edition): Chibueze Ebii
English editor: Caspar Shaller
Proof reader: Pia Hartmer, Judith Höppner, Joan Lanfranco, Heinrich-Böll-Stiftung European Union

Contributors: Johanna Bär, Ulricke Bickel, Silke Bollmohr, Larissa Mies Bombardi, Clara Bourgin,
Wolfgang Bödeker, Carsten Brühl, Helmut Butscher-Schaden, Christine Chemnitz, Henrike von der Decken,
Sunday Ekesi, Dave Goulson, Benjamin Gräub, Susan Haffman, Johannes Heimrath, Carla Hoinkes,
Heike Holdinghausen, Catherine Kunyanga, Dominic Lemken, Layla Liebetrau, Jochen Luckscheiter, Martha Mertens, Samira Mohamed,
Moritz Nabel, Saliou Niassy, Ikenna-Donald Ofoegbu, Cecilia Moraa Onyango, Andre Prescher, Ilang-Ilang Quijano, Anna Satzger,
Achim Spiller, Sevgan Subramanian, Chrysantus Tanga, Lisa Tostado, Katrin Wenz, Johann Zaller, Anke Zühlsdorf

The views do not necessarily reflect those of all partner organizations.
The maps show the areas where data are collected and do not make any statement about political affiliation.

Cover image: © Martina Puchalla, STOCKMAR+WALTER Kommunikationsdesign

Editorial responsibility (V. i. S. d. P.): Annette Maennel, Heinrich-Böll-Stiftung

1st edition, November 2022

ISBN: 978-9-46400747-3
Legal Deposit: D/2022/11.850/2

This material – except the cover image, publication covers and logos – is licensed under the Creative Commons “Attribution 4.0 International” (CC BY 4.0). For the licence agreement, see <https://creativecommons.org/licenses/by/4.0/legalcode>, and a summary (not a substitute) at <https://creativecommons.org/licenses/by/4.0/deed.en>. Individual graphics from this atlas may be reproduced if the attribution “Eimermacher/stockmarpluswalter, CC BY 4.0” is placed next to the graphic. Please cite as: “Pesticide Atlas 2022, Heinrich-Böll-Stiftung & others”



Friends of the Earth Europe gratefully acknowledges financial assistance from the European Commission (LIFE Programme). The sole responsibility for the content of this document lies with Friends of the Earth Europe. It does not necessarily reflect the opinion of the funder mentioned above. The funder cannot be held responsible for any use that may be made of the information contained therein.



FOR DOWNLOADS

Heinrich-Böll-Stiftung Abuja, Nigeria
<https://ng.boell.org/en/pesticides-atlas-nigeria>



PESTICIDE ATLAS

Facts and figures about toxic chemicals in agriculture

NIGERIA EDITION
2022

TABLE OF CONTENTS

02 IMPRINT

06 INTRODUCTION

08 TWELVE BRIEF LESSONS ABOUT PESTICIDES IN AGRICULTURE

10 PESTICIDES AND AGRICULTURE DANGEROUS SUBSTANCES

12 CORPORATIONS BIG PROFITS WITH TOXIC TRADE

14 PESTICIDE USE IN AFRICA INCREASINGLY VULNERABLE

16 NIGERIA TIME FOR A DETOX IN AGRICULTURE

18 APPROVAL PROCEDURES UNDERESTIMATED RISKS

Parts of global crop production are lost to pests and plant pathogens each year. Pesticides have been designed to prevent these yield losses – but they also give rise to new problems.

The global pesticide market is growing – and there are only a few corporations that are dividing it up among themselves. They are increasingly investing in countries in the Global South, where pesticides are less strictly regulated.

Pesticide imports into African countries are increasing. However, there is not enough information about how they are used and the impacts they are having on human health and the environment. Governments must increase investments into alternative pest control strategies, like Integrated Pest Management (IPM).

As one of the largest importers of pesticides on the African continent, Nigeria faces mounting challenges due to their high use. A significant number of these pesticides are hazardous and already banned in regions like the European Union.

Before they are put on the market, pesticides go through an approval process in which their impacts on human health and the environment are tested. But their indirect effects on food chains and biodiversity receive little attention, neither do the effects of pesticide mixtures that are hard to predict.

20 HEALTH SEVERE CONSEQUENCES

22 SOILS IMPACT ON THE INVISIBLE ECOSYSTEM

24 RESIDUES IN KENYA TOXIC SIDE DISH

26 BIODIVERSITY EXTINCTION IN FULL SWING

28 BENEFICIAL INSECTS SMALL CREATURES WITH BIG IMPACTS

30 WATER GO WITH THE FLOW

385 million people fall ill every year from pesticide poisoning. The United Nations intend to improve the worldwide handling of pesticides to prevent harm, but there is little effective legal regulation.

Insufficient attention is being paid to pesticides accumulating in the soil, where they exert direct and indirect adverse effects on soil life – sometimes for decades.

Pesticide use leads to residues in food to which many people around the world are exposed. Kenya is no exception. Global trade of pesticide products and produce means that contaminated food is every consumer’s responsibility.

Experts have been warning for years that biodiversity is at stake. Pesticides have been identified as one of the causes for why the abundance of animals and plant species is deteriorating so quickly and disastrously.

Beneficial insects provide enormous ecosystem services to Africa. However, to continue benefitting from these natural processes, sustainable farming and environmental management practices need to be promoted.

Studies regularly reveal how pesticides contaminate rivers, lakes, coastal waters and groundwater. The pollutants often originate in agriculture and enter surface water via infiltration, surface runoff and drift.

32 LONG-RANGE TRANSPORT GONE WITH THE WIND

34 INSECT DECLINE AN ECOLOGICAL ARMAGEDDON

36 HIGHLY HAZARDOUS PESTICIDES A GLOBAL HUMAN RIGHTS CONCERN

38 GENETIC ENGINEERING MODIFIED CROPS, MORE PESTICIDES

40 SMALLHOLDER FARMERS NEW MARKETS, LESS REGULATION

42 IMPORTS AND EXPORTS BANNED BUT SOLD ANYWAY

Pesticides rarely stay in the place where they have been applied. Wind can move dust, particles, and droplets to residential areas close to agricultural land – or carry it to places many kilometres away. Approval processes are largely ignoring this problem.

Insects provide pollination services to flowering plants, control pests and ensure abundant harvests. For a long time their populations have declined dramatically – causing disaster for humans and nature. Pesticides are considered a major reason for the decline.

Substances that are proven to present a particularly high level of acute or chronic risk to health or the environment are commonly referred to as Highly Hazardous Pesticides (HHPs). Far too rarely are these substances withdrawn from circulation – especially in the Global South they cause great harm.

Genetically modified crops were supposed to reduce the use of chemicals in agriculture, reduce workloads, and increase crop yields. These promises could not be kept.

In Africa, fewer pesticides are used than in other regions of the world. Nevertheless the 33 million smallholders are increasingly becoming the focus of pesticide companies. There they also sell what has been banned in the European Union.

Many pesticides are banned in the European Union. It is illegal to use them in EU Member States, yet it is allowed to produce and export them to third countries – where they pose great risks to people and their environment.

44 GLYPHOSATE A MANUFACTURED CONTROVERSY

46 GENDER AT THE FOREFRONT OF EXPOSURE

48 DIGITAL TRANSFORMATION WHO REALLY BENEFITS FROM DIGITALIZATION?

50 EU POLICIES OBJECTIVES ARE NOT ENOUGH

52 PESTICIDE-FREE REGIONS GOOD EXAMPLES

54 AUTHORS AND SOURCES FOR DATA AND GRAPHICS

Bayer and other companies are fighting for the re-approval of glyphosate in the EU. To do so, they must prove that their pesticide active substance is not carcinogenic. But the studies presented are old – and point to the opposite.

Women working in agriculture often have lower levels of income and lack decision-making power. There is urgent need for gender equality to achieve food security and protection from pesticide exposure.

Agricultural robotics, drones, and algorithm-driven technologies for a new way of farming are becoming big business. They are supposed to help farms cut pesticide use, but there are serious doubts whether this will work.

Even though the EU’s pesticide legislation is the most sophisticated, it has failed to achieve a reduction in use. The EU’s Farm to Fork Strategy seeks to change that. Many questions remain.

All over the world, projects are proving that an ecological future is possible: More and more cities, states, and regions are trying to slash pesticide use; or even ban chemical agents completely from their fields and land.

INTRODUCTION

Traces of pesticides from agriculture can be detected everywhere – in beer and in honey, on fruit and on vegetables, in grass on playgrounds and even in urine and in the air. We have known for a long time now that pesticides have a negative effect on human health. We have also known that they cause severe harm to insects and plants and contaminate water bodies for years.

Rachel Carson published her globally acclaimed book “Silent Spring” as early as 1962, in which the biologist described the harmful effects of pesticide use. Her book is considered groundbreaking for the environmental movement and one of the most influential non-fiction books of the 20th century. Since then, many pesticides have been taken off the market. New ones have been introduced with the promise that they are less harmful to people and the environment. Promises that have rarely been kept.

Sixty years after Carson’s book came out, greater amounts of pesticides are being applied worldwide than ever before, despite stricter regulations in approval processes, and voluntary as well as binding agreements on the handling of pesticides. According to a recent study, the number of people affected by pesticide poisoning each year has risen to 385 million, and pesticide use is considered a major cause of species decline. Especially in countries with high levels of biodiversity, the cultivation of genetically modified soy has contributed to a dramatic increase in pesticide use.

“**Never in history have pesticides been used so pervasively**

Hardly any country in the world has an ambitious pesticide reduction strategy or even ideas for an agriculture that is truly independent of chemical crop protection. There is a reason for this: The market for pesticides is highly lucrative. Only a few well-connected and influential agrochemical companies divide it up among themselves. At the forefront: German companies like Bayer and BASF. No wonder, then, that Germany is one of the largest pesticide exporters of the world.

The promising growth markets of pesticide companies have long since ceased to be in Europe and are mainly found in Latin America and Asia today. But pesticides are also increasingly being exported to African countries. Nigeria is one of the largest importers of pesticides on the African continent. It already faces mounting public health and environmental challenges due to their high use. A significant number of these pesticides are no longer authorised or banned in the EU due to their harmful effects.

A key demand of Nigerian civil society is therefore to create regulations and laws that ban these toxic substances from being imported. The fact that the new German government has committed

itself to create laws that will prohibit their export is a welcome development.

However, an import ban on the most toxic substances does not go far enough. There has to be more thorough registration processes, effective controls and, at the same time, support for farmers to use fewer pesticides. Alternative agricultural strategies like integrated pest management, the use of biopesticides, organic farming and agroecology are still largely overlooked. Like the EU, Nigeria should chart a path towards an overall reduction in pesticide use.

“**An ecological turnaround requires an agricultural change**

The climate crisis will increase the spread of plant diseases, pest infestations, and extreme weather in many parts of the world. To reduce the increased pressure on beneficial and essential insect and plant populations, our agricultural systems must adapt to meet these challenges with fewer pesticides. To do so, they must become more diverse and protect and use beneficial insects as allies. We face an urgent task: To learn how to fight alongside nature instead of against nature. To succeed, we must set the course now. That is why we want to use this atlas to provide data and facts to facilitate a lively debate.

Jan Philipp Albrecht and Imme Scholz
Heinrich-Böll-Stiftung

Jochen Lukscheiter
Heinrich-Böll-Stiftung, Abuja Office

Olaf Bandt
Bund für Umwelt und Naturschutz Deutschland

Doris Günther
Pestizid Aktions-Netzwerk

12 BRIEF LESSONS

ABOUT PESTICIDES IN AGRICULTURE


1 The global consumption of pesticides is increasing, even though the harmful health and environmental effects are known. International goals of **BIODIVERSITY CONSERVATION** can only be achieved if the use of pesticides is significantly reduced.

2 Herbicides are applied against unwanted plants and are the **MOST USED GROUP OF ACTIVE SUBSTANCES**. Insecticides are effective against harmful insects. Often even in smallest amounts and even against other insects that were not targeted.



3 About 385 million cases of **PESTICIDE POISONING** occur worldwide every year. In Nigeria, pesticide poisonings are a regular occurrence but injuries and deaths are not systematically captured by government agencies.

7 Pesticides **CONTAMINATE** water via infiltration, surface runoff and drift. They also accumulate in the soil and exert adverse effects on soil life – sometimes for decades.



8 Pesticide residues in food can be **HARMFUL TO PEOPLE'S HEALTH**. In June 2015, the European Union banned the import of dried beans and other Nigerian agricultural products that contained levels of pesticide residues considered dangerous to human health.



9 Four corporations from the Global North control 70% of the global pesticide market. They are **EXPANDING THEIR BUSINESS** to low- and middle-income countries where pesticides are less strictly regulated.



4 Pesticides that are **NOT ALLOWED FOR USE** in countries where they are produced are still exported to other countries. In Nigeria, more than half of all the pesticide products registered have been withdrawn from the European market or are heavily restricted.



5 Nigeria's pesticide registration process involves the risk-based evaluation of comprehensive data. However, the harmful effects of **PESTICIDES ON WHOLE ECOSYSTEMS AND THE LOCAL CONTEXT** are not systematically taken into account when pesticides are approved for use in Nigeria.



6 Pesticide active ingredients usually do not stay in the place they were applied. They can seep into the soil and **GROUNDWATER**, become airborne, or blow away – some can be found over 1,000 kilometres away from the point of application.



10 Pest populations in Africa are kept in check by natural enemies like dragonflies, predatory bugs and beetles. Unsustainable and intensive farming practices such as use of pesticides, monocultures and land use changes disrupts this natural balance. **PROTECTING ENVIRONMENTS FOR BENEFICIAL INSECTS** is important for Nigeria's agricultural biodiversity.



11 United Nations experts have considered highly hazardous pesticides (HHPs) a **GLOBAL HUMAN RIGHTS CONCERN** for a long time. Pesticides endanger among other things, the right to food, health and a clean environment. Nigerian civil society organisations are therefore calling on the government to do more to reduce the use of pesticides and ban particularly toxic ones.



12 Unlike industrial monocultures, agroecological cultivation practices, including more crop rotations and combinations, enable farmers to use less or no pesticides. This is also more economically viable for smallholders. Some regions of the world are going ahead. But a binding international **TREATY ON THE REDUCTION OF PESTICIDES** does not yet exist.

PESTICIDES AND AGRICULTURE

DANGEROUS SUBSTANCES


Parts of global crop production are lost to pests and plant pathogens each year. Pesticides have been designed to prevent these yield losses – but they also give rise to new problems.

Grave famines and economic upheavals resulting from crop failures have occurred throughout history. People have always fought against this existential challenge—for example by using certain cultivation methods and certain crop rotations to avoid weeds and pests. The industrial revolution saw the emergence of the first synthetic chemical pesticides: They were meant to protect crops and reduce workloads. Starting in the 1940s, the chemical industry began marketing broad-spectrum pesticides – they were poisonous to entire groups of organisms and initially proved to be much more effective compared to previously available substances. Global pesticide use has continued to grow steadily for decades: Between 1990 and 2017 by about 80 percent. The interplay of pesticides, fertilizers and technological progress led to a fundamental change of agricultural production. As farmers now kept diseases and pests at bay through pesticides rather than crop rotations and crop combinations, monocultures of single crops repeatedly grown on the same land became the standard. As a result, today’s industrial agriculture is dependent on pesticides and is largely unimaginable without them. Capital-intensive inputs increased yields in many industrialized countries since the 1950s. Therefore, the supply of agricultural products grew much faster than the demand; a development that has resulted in lower prices for agricultural products, which become cheaper and cheaper, while wages for farmers and agricultural workers have decreased. Not only has the amount of pesticides applied worldwide increased, but so has the scientific research on pesticide effects – experts have gained more and more knowledge about how pesticides can affect human health and pollute the environment.


Today, pesticide consumption worldwide stands at 4 million tonnes globally. Half of the substances applied are herbicides, which are used against weeds; about 30 percent are insecticides, which are used against insects that can harm harvests. And about 17 percent are fungicides against fungal infestation. The global pesticides market size reached a value of nearly 84.5 billion US dollars in 2019, with an annual growth rate of more than 4 percent since 2015. In the next few years, the rate of growth could increase further. By 2023, the total value of all pesticides used is expected to grow at a rate of 11.5 percent to nearly 130.7 billion US dollars. Many factors, like soil degradation and biodiversity loss, have contributed to the increase. The climate crisis can be another driver for pesticide use. A study from the US-American Seattle University found: Insect activity in crop-growing regions will rise along with temperatures. This will boost losses of rice, maize and wheat by 10–25 percent for each degree Celsius that temperatures rise. There are major reasons for this. For example, climate crisis is altering pest populations and the ratio of pests to beneficial insects. Insects seek out conditions that suit them and move to new areas that lack their natural enemies. This will cause their populations to grow, resulting in more crop damage. Furthermore, the plants’ natural potential to resist to pests decreases as a result of climate-related stress.

Depending on the region and the phase of industrial development, usage of pesticides is associated with different intensity. The 1960s are considered the age of the “Green Revolution” that was devised to increase agricultural production, particularly in the Global South – through the use of pesticides, fertilizers, high-yield crops and irrigation. In retrospect, civil society organizations and scientists view the “Green Revolution” as the beginning of a failed agricultural development, which led many farmers into desperate situations.

A small number of corporations from the Global North divide the multibillion dollar market between themselves



NO REDUCTION TO BE SEEN
Pesticide use in tonnes by continent in 2020 and change since 1999




Many people in the Global South have gone into debt to buy expensive means of production. Due to high profit margins and insufficient government regulation, the trade in illicit pesticides has increased over recent years. And the sale of counterfeit pesticides has become a profitable business as well: In the first four months of 2020, illegal pesticides worth up to 94 million euros were seized in the EU and six other non-EU countries such as Colombia, Switzerland and the USA. The use of such pesticides puts farmers at particular risk because the ingredients and their concentrations may be misstated or misrepresented – making their effects and toxicity unpredictable.

Pesticides do not stay where they have been applied. They contaminate the environment and contribute to an imbalance in the ecosystem. New research shows that pesticides even contribute to pollution with microplastics when active ingredients are intentionally encapsulated for slower release. A key challenge for governments is to inform farmers worldwide about the dangers of pesticides, to take measures to protect them and to enable manageable crop protection alternatives to chemical pest control. Ideas on how this could work around, although research in topics such as ecologically-based pest management remains underfunded. ●

Neonicotinoids are applied to fields at lower doses than conventional pesticides, but are highly toxic. They have led to annual rates of reductions in insectivorous birds by 3 percent

The global pesticide market is growing. South America and Africa are among the markets with the highest growth rates – but differ largely in current use and application rate



BIG PROFITS WITH TOXIC TRADE

The global pesticide market is growing – and there are only a few corporations that are dividing it up among themselves. They are increasingly investing in countries in the Global South, where pesticides are less strictly regulated.

Agrochemical companies such as Bayer or Syngenta emerged from chemical or pharmaceutical companies – some of which were founded already in the 19th century. In the mid-1990s, with the advent of genetic engineering in agriculture, they discovered a new business model: combining pesticide sales with seed sales. In order to form new specialized groups, they bought up smaller seed producers in large numbers and, around the turn of the millennium, split off the agricultural division from the rest of the business. In recent years, the shares of these corporations in the global market have increased sharply once again. In 2015, the US corporation Dow Chemicals had announced a merger with Dupont. Both companies combined their pesticide and seed businesses to Corteva Agriscience four years later. In 2017, the Chinese state-owned enterprise ChemChina took over the Swiss agricultural group Syngenta. In 2018, the German chemical company Bayer acquired the U.S. company Monsanto and sold parts of its business to German chemical company BASF, which entered the seed business with the acquisition. And in 2020, Syngenta, the Israeli pesticide company Adama, and Sinochem from China were combined to form Syngenta Group.

The top four firms – Syngenta Group, Bayer, Corteva and BASF – controlled around 70 percent of the global pesticide market in 2018. 25 years earlier, their market share was only 29 percent. In the seeds sector – now led

by exactly the same groups – the share of the biggest four rose from 21 to 57 percent over the same period.


The power of these players and the continued merging of the two business models have implications for product range and agriculture worldwide: Pesticide selling seed producers have an interest in ensuring that their agrochemicals are also used in the cultivation of their seed. The leading global providers of seeds and pesticides focus on selective breeding and genetic modification of a small number of crops. First and foremost, soybean and maize. They account for about two-thirds of the seed market's volume. Bayer generates about 75 percent of its seed sales from maize and soybeans, Syngenta 55 percent and Corteva a full 85 percent.

Aiming to further developing seeds, the big companies have increased their research expenditures in recent years, while research expenditure in the agrochemical sector has been stagnating at the same time. In 2000, 70 percent of global agrochemical sales were patented or proprietary formulations. Since then, patents on popular agrochemicals have expired, with no new patented active ingredients to take their position on the market. Meanwhile only 15 percent are patented. One reason for this can be found in stricter approval procedures, largely in the European Union – which led to an increase in cost for bringing a new active ingredient to market. In light of these costs, major firms tend to use older active ingredients, combined in new mixtures.

The HHP list of PAN International currently contains 338 highly hazardous pesticides with high levels of acute or chronic hazards to health or environment according to internationally accepted classification


TOXIC TOPSELLER

The bestselling Highly Hazardous Pesticides (HHPs) in 2018, by company



HIGHLY HAZARDOUS AND HIGHLY PROFITABLE

Percentage of Highly Hazardous Pesticides (HHPs) of the five biggest pesticide companies' total revenue, and highly hazardous substances' turnover on the five most important markets in 2018, in million US dollars



The best-selling pesticide products include the herbicide glyphosate (patented in 1971, on the market since 1974), paraquat (herbicidal effect discovered in 1955, on the market since 1962), the herbicide atrazine (on the market since 1958) and neonicotinoids, a new class of insecticides (on the market since the early nineties). What they all have in common is that they are considered dangerous: Glyphosate for example is suspected of being carcinogenic, paraquat is highly toxic to humans, atrazine is hormone-disrupting and neonicotinoids are highly toxic to bees.

In industrialized countries, the five largest producers sell less highly dangerous pesticides overall than in Asia, Africa and Latin America: While they account for 12 percent of total pesticide sales in Germany and 11 percent in France, they account in Brazil for 49 percent and in India for 59 percent. One reason for this is that the EU and the countries of the European Free Trade Association (EFTA) have banned several Highly Hazardous Pesticides. Elsewhere, however, these substances are still permitted due to incomplete regulation – especially in South America, Asia, and Africa, where pesticide sales are on the rise.

The continuous growth of the global pesticide market by an average of 4 percent annually is mainly due to sales in these world regions. Africa still uses the least pesticides, with an average of less than 0.4 kilograms per hectare of cropland, while


In the last 30 years, the value of pesticide exports from the EU has multiplied. Highly Hazardous Pesticides (HHPs) which account for about one third of the more than 1,000 active ingredients worldwide, are also amongst them

European companies issued plans in 2018 to export 81,000 tonnes of pesticides prohibited on their own fields. Main destination: the Global South

worldwide the figure is around 2.6 kilograms per hectare but is starting to catch up with other regions: Industry has long since identified the African continent as its largest growth market. With the increasing presence of the agricultural industry, the use of Highly Hazardous Pesticides is also increasing. ●

SOLD, SHIPPED, POISONED

Pesticide export value in 2020, European Union



PESTICIDE USE IN AFRICA

INCREASINGLY VULNERABLE

Pesticide imports into African countries are increasing. However, there is not enough information about how they are used and the impacts they are having on human health and the environment. Governments must increase investments into alternative pest control strategies, like Integrated Pest Management (IPM).

Africa is quickly developing a dependency on agricultural pesticides. Increasing farm labor costs and pressure to intensify land productivity as well as greater availability of internationally manufactured, inexpensive products are some of the factors driving increased use on the continent. However, most African governments do not have adequate resources to monitor the impacts of pesticides or the capacity to prevent negative human health and environmental consequences. The widespread adoption of pesticides means that millions of smallholder farmers are exposed to the risks associated with using these chemicals on their farms.

Over the last five years, pesticide imports into Africa have increased significantly. In West Africa the imports have doubled in five years from 218,948 tonnes in 2015 to 437,930 tonnes in 2020. In 2020, Nigeria's imports alone (147,446 tonnes) exceeded the total imports of Southern Africa (87,403 tonnes) and North Africa (109,561 tonnes). Despite increasing imports in these regions, the informal nature of agricultural production has made it difficult to record how pesticides are used hence the big differences between the imported quantities and use data. For example, in 2020 the FAO recorded the value of imports in Southern Africa at 87,403 tonnes, compared to 27,000 tonnes of pesticides that were used.

New traders and business people with limited knowledge about pesticides are taking advantage of the situation. There has been a proliferation in the illegal trade of counterfeit products which are smuggled through porous borders. Counterfeit pesticides are not authorised for sale by the mandated pesticide authorities and can lead to a total loss of treated crops, compromising farmer livelihood. In addition, counterfeit pesticides might contain chemicals that are either banned or restricted. Regulators are usually caught unaware when produce from their countries includes active ingredients that they have not approved.


According to the FAO pesticide use data from 2020 and in terms of leading countries in each region, South Africa leads in Southern Africa (26,857 tonnes), Egypt in North Africa (11,352 tonnes), Cameroon in West Africa (7,322 tonnes) and Ethiopia in Eastern Africa (4,128 tonnes).

Despite ever increasing pesticides use, the prevalence of food insecurity and malnourishment is not improving. Between 2019 and 2021 approximately 20 percent of people on the continent were undernourished, which has increased from 16 percent between 2016 and 2018.

There are differences in the quantities and types of pesticides used within and between the regions.


For example, while Egypt uses more fungicides than herbicides, Ethiopia uses more herbicides than fungicides. And, although Cameroon and Ghana are in the same region, Ghana uses more herbicides (6,498 tonnes) than Cameroon (4,801 tonnes). These differences can be attributed to the type of production that characterizes the country's ag-

Despite increasing pesticides imports in Africa, data on pesticides use is lacking. The information we don't have is reason for concern, because the impact of pesticides on the environment, farmers and consumers is likely underestimated



MORE PESTICIDES, NOT LESS HUNGER

Top ten highest importers of pesticides in Africa, 2020, in tonnes



gricultural model. The data provides important information about the pest problems facing farmers and therefore the ecology specific solutions that can be used in varying local contexts.

An effective management strategy for pesticides in Africa must include educational programs focusing on pesticide use, health and environmental risks, stringent policies on imports of highly hazardous pesticides (HHPs), increased government support to production systems that promote agroecology and long-term monitoring on the impacts of pesticides.

African countries are party to many conventions on the protection of human health and the environment such as Stockholm Convention on Persistent Organic Pollutants, the Montreal Protocol and the International Code of Conduct on the Distribution and Use of Pesticides. The Inter-African Phytosanitary Council is a specialized technical office of the African Union mandated to among other things, harmonise pesticides registration on the continent. Regional institutions responsible for ensuring the safe use of pesticides also exist and include for example, the West Africa Committee for Pesticide Registration. In addition, each of the 54 African states has its own system that governs pesticide registration

Companies argue that synthetic inputs will increase crop production to ensure food security and nutrition. However, the prevalence of undernourishment in Africa is steadily rising despite increasing pesticide imports and use

and use. It is worrying that despite the institutional frameworks for judicious pesticides management, most countries in Africa have not implemented an effective impact monitoring process and investments into farmer training on alternatives are lacking.

Civil society organisations advocating for the adoption of ecologically sustainable food systems and a ban on HHPs, fill a gap that governments are failing to address. Organisations such as the Pesticide Action Network-Africa which is based in Dakar, for example, are working to eliminate the harm caused by pesticides. Other organisations such as the Organic Producers and Processors Association of Zambia, Kenya Organic Agriculture Network, Egypt Biodynamic Association, and the National Federation for Organic Farming in Senegal provide training to promote IPM strategies whilst sensitizing producers, handlers, processors, consumers and policymakers on the negative effects of pesticide use on human health and the environment. ●

TIME FOR A DETOX IN AGRICULTURE

As one of the largest importers of pesticides on the African continent, Nigeria faces mounting challenges due to their high use. A significant number of these pesticides are hazardous and already banned in regions like the European Union.

Nigeria is well known for its oil wealth, but the agricultural sector is the backbone of the country's economy. In 2021, agriculture accounted for about 26 percent of the gross domestic product, compared to oil and gas (7 percent), manufacturing (9 percent) and trade (16 percent). About 35 percent of the active labour force is directly employed in agriculture.


However, the sector faces major challenges. It needs to feed a rapidly expanding population in a warming climate characterised by increasingly frequent floods and droughts that reduce soil fertility, while parts of the country remain in conflict, ranging from the Boko Haram crisis in the northeast to farmer-herder clashes in the Middle Belt. The ongoing Covid-19 pandemic and the impacts of the war in Ukraine on global energy prices have further exacerbated food price inflation.

The country is currently unable to meet its domestic requirements for food as population growth has outpaced the rate of food production. According to the United Nations,


the number of people living in Nigeria is expected to almost double by 2050 from its current level of just over 200 million, fuelling greater consumer demand for food. Government policies have therefore focused on increasing productivity. These mainly promote conventional agriculture that requires a high degree of external inputs, such as pesticides and artificial fertilisers. However, the use of pesticides has resulted in negative health, environmental and economic consequences in Nigeria.

The registration of pesticides in Nigeria is covered by the Drugs and Related Products Act (1996), the Pesticide Registration Regulations (2019) and the Guidelines for Pesticide Registration. The National Agency for Food and Drug Administration and Control (NAFDAC) is the registration body and also regulates and controls the import, export, manufacture, advertisement, distribution, sale and use of chemicals. As at November 2022, NAFDAC's Green Book product database lists 682 synthetic chemical pesticide products (excluding chemical repellents). More than half of these products include active ingredients that are not approved in the European market due to, for example, their potential chronic health effects, environmental persistence, high toxicity

There is no reliable data over time for Nigeria's pesticide imports, but the latest information from the Food and Agricultural Organisation (FAO) shows that the country is the largest importer on the continent by volume (see p. 15), spending \$384 million in 2018



BANNED YET ACCEPTED
Products registered in Nigeria and not approved in Europe



for fish or bees, or insufficient data to uphold the principle of preventing harm. NAFDAC is reviewing registered active ingredients with the intention to phase out some of the most hazardous substances – including ametryn, chlorpyrifos and methomyl – and to reclassify the use of a few others. The outcomes of this process are yet to be published.

Surveys have shown that 80 percent of the pesticides used most frequently by small-scale farmers are highly hazardous pesticides. Among the most commonly used are atrazine, chlorpyrifos and mancozeb – all of which are prohibited in the European Union.

Nigeria has missed important export opportunities because of the high use of pesticides on farms and during storage. In June 2015, the European Union banned the import of dried beans and other Nigerian agricultural products that contained levels of pesticide residues considered dangerous to human health. The United States has similar bans in place.

The use of these substances also impacts the health of the Nigerian population. Although the Nigerian government does not systematically collect data on pesticide poisonings, there are regular media reports of such incidents. In 2020, more than 270 people died in Benue State after drinking river water poisoned with Endosulfan, a pesticide that was banned globally in 2011 under the Stockholm Convention, including in Nigeria, but is still available in Nigerian markets.

Part of the reason why these challenges persist is the lack of interaction and collaboration between the range of agencies involved in the pesticide registration, regulation and enforcement system. These include the following, among others: the Nigeria Agricultural Quarantine Services (NAQS), under the Federal Ministry of Agriculture and Rural Development, is responsible for residue monitoring for export certification and for pest monitoring. The Standards Organization of Nigeria, under the Federal Ministry of Industry,

More than half of all the pesticide products registered in Nigeria are not approved in the European Union due to their toxicity

Trade and Investment, develops and enforces standards for laboratory pesticide analysis. The National Environmental Standards and Regulations Enforcement Agency, under the Federal Ministry of Environment, enforces regulations. The Department of Food and Drugs Services, under the Federal Ministry of Health, develops policies, guidelines and programmes to ensure food safety in Nigeria.

Every one of these agencies works in a silo with insufficient communication between role players. There is no dedicated legislation to govern the sector, but a National Pesticide Policy, led by NAQS, was announced in March 2019. This is intended to combine all the related responsibilities, policies and guidelines, but no further progress has been published. In addition, the lack of distinct legislated registration pathways for biopesticides/biological-control agents hinders the implementation of alternative agriculture.

Sustainable agricultural techniques like organic farming or agroecology have not yet appeared at significant scale in Nigeria. The national agricultural budget for 2021 allocated 60 percent of the total capital budget to research and development, but only 1.3 percent to sustainable agricultural techniques. With no top-down policies on these strategies, the movements for sustainable agriculture are mostly driven from below by NGOs and farmers.

However, some changes are on the horizon. The Nigerian government's recently published National Agricultural Technology and Innovation Policy (2022–2027) identifies the need for a regulatory framework for the adequate control and monitoring of the pesticide industry as well as for the development of an organic agriculture policy. ●

APPROVAL PROCEDURES

UNDERESTIMATED RISKS

Before they are put on the market, pesticides go through an approval process in which their impacts on human health and the environment are tested. But their indirect effects on food chains and biodiversity receive little attention, neither do the effects of pesticide mixtures that are hard to predict.

EU approval of pesticides is carried out in a two-stage process overseen by the European Food Safety Authority (EFSA). In the first step, active ingredients are approved at the European level, which is divided into different geographic areas: EFSA distinguishes three European zones with comparable ecological and climatic conditions, namely North, Central and South. In the second step, the pesticide products containing these active ingredients are approved by individual EU Member States. The pesticide manufacturers submit their scientific information and studies at the EU level which provides the data necessary to conduct the environmental and health risk assessment. EFSA then commissions different Member States – appointed as rapporteurs – to review these dossiers. The rapporteur prepares a Draft Assessment Report with regards to the risks for humans and the environment which EFSA peer reviews, together with the Member States. If this process concludes that there are no unacceptable effects on environment and human health to fear, the agency gives approval. This ultimately



means that adverse effects on the environment or on non-target organisms can not stop registration if they are considered acceptable. This may occur, for example, if a beneficial insect population of for instance ladybugs was to recover after pesticide application.

During the review process, EFSA works with the EU Commission and Member States, and carries out public consultations which includes stakeholder surveys designed to collect the views of stakeholder organizations and Member State Authorities. EFSA prepares a final draft report and a committee of Member State representatives votes on the draft decision. The decision on whether to approve the substance is taken by the European Commission in consultation with Member States.

The approval of an active ingredient is granted for a defined number of years, not exceeding 10 years. For a renewal new data must be included in the decision-making process. It is important to note that active ingredients which meet with certain cut-off criteria – a classification as mutagenic, carcinogenic or harmful to reproduction and endocrine system – will not be approved in the EU.

Despite independent studies suggesting otherwise, the herbicide glyphosate was granted re-approval by the EU in 2017. The controversial herbicide was first approved in 2002

The results from approval tests with only a few species are subject to uncertainties. To compensate for these uncertainties, safety factors are supposed to help



under the new EU pesticide legislation. Previously, it was only permitted in some Member States. The re-registration of glyphosate was scheduled for 2013, and Germany served as rapporteur country, with Slovakia as co-rapporteur. The process received widespread attention due to environmental and health concerns; meanwhile, the International Agency for Research on Cancer (IARC), which as part of the World Health Organization (WHO) has devised a system of categories to evaluate the carcinogenicity of a substance to humans, has classified glyphosate as “probably carcinogenic” to humans. However so far only Luxembourg was the first EU country to ban glyphosate. However so far only Luxembourg was the first EU country to ban glyphosate. The main reason for differing assessments was that the IARC used independent studies for evaluation, while the national regulatory authorities relied on manufacturer studies. Furthermore, the IARC assessed glyphosate containing products and occupational exposure, while national authorities mainly considered the pure active ingredients only, dietary exposures and risks to the general population. As a compromise, the approval of glyphosate was only granted for another five years instead of ten years. An alliance of glyphosate manufacturers called Glyphosate Renewal Group (GRG) has already submitted a dossier to EFSA to ensure that the herbicide continues to be approved after 2022. It comprises 180,000 pages. To address this, the Commission appointed four Member States acting

Dangerous pesticides must be phase out. Biopesticides can be an option for substitution if other measures within the framework of integrated pest management have failed.

jointly as ‘rapporteurs’, known as the Assessment Group on Glyphosate (AGG), consisting of EU Member States France, Hungary, the Netherlands, and Sweden.

Although pesticides must meet the strict EU approval criteria, the current environmental impact assessment does not seem to prevent the approval of pesticides that have harmful effects on the environment. The EFSA guidelines focus on how to evaluate the impact of active ingredients with consideration to surrogate species of birds, mammals, honeybees, wild bees or earthworms. Ecologists and civil society organizations demand that the impacts on fungi, amphibians, bats, reptiles, or wild plants are also considered. Interactions between organisms and indirect pesticide effects are left out the approval process as well. Another important aspect not considered in environmental risk assessments is the fact that most agricultural crops are treated not only with a single pesticide but with a variety of pesticides each season. These mixtures’ environmental effects are still largely unknown – evidence is mounting that they are stronger than the effects of individual substances. Because of these fundamental flaws pesticides can not be considered safe for the environment. ●

HEALTH

SEVERE CONSEQUENCES

385 million people fall ill every year from pesticide poisoning. The United Nations intend to improve the worldwide handling of pesticides to prevent harm, but there is little effective legal regulation.

People can be unintentionally exposed to pesticides in various situations: on the field, in the forest, through food or drinking water. The clinical diagnosis of pesticide poisoning is made when typical symptoms develop after exposure. Some health effects may occur right away, while other symptoms may occur several hours after exposure. Short-term adverse health effects are called acute effects, including stinging eyes or rashes. The victim may feel tired and listless and suffer from headaches and aching limbs. The digestive tract is also frequently affected – the

consequences are nausea, vomiting or diarrhoea. In serious cases of poisoning, the victim’s organs can fail: the heart, lungs or kidneys stop functioning. The total number of fatalities around the world from unintended pesticide poisonings are estimated at some 11,000 per year.

Farmers are at a higher risk of getting exposed to pesticides, but the substances can also pose risks to people outside the agricultural sector as pesticides are mobile and difficult to control. They often contaminate the environment and end up in our food.

The lack or misregarding of safety precautions can result in serious injuries or fatalities as the following two examples show: In 2013, twenty-three school students in Bihar, India, died within minutes of eating a meal of rice and potato curry that was part of a lunch program against malnutrition. The forensic investigation found that the meal had been prepared with cooking oil that contained the pesticide monocrotophos. In the same year, an airplane sprayed an insecticide over a rural school in the Rio Verde for a full 20 minutes. Children and their teachers were eating their lunches under the open sky when the toxic chemicals were sprayed on them. Dozens of children and adults were hospitalized. The school – located among vast maize and soy plantations – was doused in the pesticide Engeo Pleno, produced by the seed and chemical company Syngenta.

Many of those affected by poisoning suffer from long-term effects: There is a substantial body of evidence on the relationship between exposure to pesticides and elevated rate of chronic diseases such as Parkinson’s or childhood leukaemia. Pesticides have also been linked to an increased risk of liver and breast cancer, Type 2 diabetes and asthma, allergies, obesity and endocrine disorders.



Birth defects, preterm births and growth disorders can also be traced back to contact with pesticides. In recent years, a widely publicized debate has centered on glyphosate. Several people who developed cancer after being exposed to the herbicide have sued its manufacturer Bayer for damages, who has lost various lawsuits already. About 96,000 plaintiffs reached settlements estimated at 11.6 billion euros; around 30,000 of these lawsuits are still ongoing.

In March 2015, the International Agency for Research on Cancer (IARC) – an intergovernmental agency that forms part of the World Health Organization (WHO) of the United Nations – classified glyphosate as “probably carcinogenic to humans”. A 2019 University of Washington scientific meta-study found that the overall meta-relative risk of non-Hodgkin lymphoma in individuals that were exposed to glyphosate-based herbicides increased by 41 percent.

Even at low concentrations, endocrine disrupting chemicals (EDCs) are a clear health risk. For example, they are found in cosmetics, plastic packaging – or pesticides

BAD FOR BOTH HEALTH AND FINANCES

Estimated health care costs of endocrine disrupting chemicals (EDCs) in the European Union, in billion euros




Study from 2015, conservative estimates. Actual total numbers are likely to be significantly higher. Various diseases associated with EDCs such as Parkinson’s disease are not included due to lack of data

© PESTICIDE ATLAS 2022 / TRASANDE ET AL.

SUFFERING AND DYING MOSTLY OCCURS IN THE GLOBAL SOUTH

Global distribution of pesticide poisoning per year, study from 2020



Poisoning affects 44 percent of all agricultural workers worldwide – and in a low-income country like Burkina Faso as many as 83 percent

Several studies show that pesticide poisonings have been rising sharply for years – today about 385 million cases of acute poisonings occur each year. In 1990, a WHO task force estimated that about one million unintentional pesticide poisonings with severe manifestations occur annually, leading to approximately 20,000 deaths. Because many states do not have central reporting offices, it can be assumed that the actual number could be significantly higher as many cases remain unreported: Scientists point out that the total number of occupational poisonings in 1990 was even twenty-five million. One reason for the increase to 385 million poisonings today is probably the intensified pesticide use all over the globe: the worldwide tonnage increased by almost 81 percent between 1990 and 2017. This includes a 484 percent increase in South America and a 97 percent increase in Asia.

Most victims live in the Global South, where environmental, health, and safety regulations are often the weakest. The use of Highly Hazardous Pesticides (HHPs) is also

a reason for the high poisoning rate. 60 percent of deaths related to pesticide poisonings occur in India.

In order to reduce the high number of pesticide poisonings, the WHO and the Food and Agriculture Organization (FAO) – a specialized agency of the United Nations that leads international efforts to defeat hunger and improve nutrition and food security – have developed a voluntary framework and standards for pesticide management. Among other things, the code of conduct recommends avoiding pesticides that require personal protective equipment too uncomfortable or expensive to use. The guideline recommends also the use of agroecological alternatives and a ban on Highly Hazardous Pesticides (HHPs). However, these recommendations have hardly been implemented so far, they are still non-binding and without legal obligation. ●

SOILS

IMPACT ON THE INVISIBLE ECOSYSTEM

Insufficient attention is being paid to pesticides accumulating in the soil, where they exert direct and indirect adverse effects on soil life – sometimes for decades.

In healthy soils very high levels of biodiversity can be observed: Soil is home to a quarter of all known species on Earth. Soil life is so abundant that a shovelful of healthy soil contains more living organisms than there are people on Earth. It is hard to overestimate what all this teeming life in the soil is capable of achieving – tens of thousands of underground species of invertebrates, bacteria, and fungi are constantly filtering our water, recycling nutrients, counteracting soil-borne diseases, building humus, sequestering greenhouse gases, and regulating the climate. So soil is not only the substrate on which we grow our food – but also a non-renewable resource that must be treated with care.


Most pesticides are designed to be toxic to organisms and it is all the more concerning that nearly two-thirds of all agricultural land worldwide is contaminated with at least one pesticide active ingredient. In Europe, soil analyses revealed that more than 80 percent of 317 agricultural topsoils tested contained pesticide residues. The most commonly found and most highly concentrated pesticides were

the long-banned insecticide DDT, the herbicide glyphosate as well as its degradation product AMPA, and broad-spectrum fungicides such as boscalid, epoxiconazole, and tebuconazole.

Pesticide residues in the soil affect soil life. A systematic review of nearly 400 published studies found: Pesticides harm organisms that are vital for maintaining healthy soils in over 70 percent of the more than 2,800 experiments included in this review. These effects were observed at all organismic levels: bacteria, fungi, and soil fauna. Pesticide residues in soil are also associated with the decline of earthworms, microorganisms, and symbiotic mycorrhizal fungi – which provide not only nutrients to plants but also keep them healthy.


Ecotoxicological research on pesticides has always focused specific effects, for example on how insecticides affect beneficial soil insects, or how fungicides affect soil fungi. However, pesticides have an impact that goes far beyond that: They usually have negative effects on a wide range of non-target organisms. One example is glyphosate – the most

317 agricultural topsoil samples from across the European Union were examined: Almost half contained up to 5 different residues



GOING UNDERGROUND

Median number of pesticide residues in European agricultural soils, by selected member state, study from 2019



widely used herbicide in the world. It affects soil life in a variety of ways, directly and indirectly: The use of glyphosate can harm soil bacteria and mycorrhizal symbiosis with the roots of grapes. Even 11 months after application, the herbicide can still be affecting the nutrient composition of the entire grape plant. Glyphosate herbicides reduce activity and reproduction of earthworms and can force tiny spring-tails from the soil to the surface, making them more vulnerable to predators. These impacts on soil life can further impair water infiltration after heavy rains – and lead to more glyphosate contamination in water bodies.

Pesticide use can also harm subsequent crops. Nevertheless, this is hardly taken into account in risk assessment. Persistent glyphosate residues in soil have been shown to alter many plant processes: They change the regulation of plant defence systems against diseases and harmful soil-borne fungi. Glyphosate residues in livestock feed can even be transferred to manure and affect the growth of fertilized crops the following year. Pesticides containing intentionally added microplastics also contribute to the pollution of soils. The use of such plastic-coated synthetic agrochemicals is rising, with producers marketing their controlled-release function. According to a 2019 report from the European Chemicals Agency (ECHA), microplastics added intentionally to fertilizers, pesticides and seed coatings account for nearly half of the approximately 51.500 tonnes of microplastics used each year in the European Economic Area.


Even after two decades of organic agriculture, up to 16 different pesticide residues were present in soil samples from 60 agricultural sites throughout Switzerland

Even years after pesticide use, the soil contamination is a problem: It has become an issue of increasing concern in Europe due to high soil persistence and toxicity to non-target species

Environmental experts are troubled by the many negative effects that pesticides have had on soil life for decades. They are calling for greater consideration of biodiversity and soil health issues when assessing the environmental risks of pesticides. In addition to common soil life, many other species also spend part of their life cycle in the soil: ground beetles, ground-nesting bees, or amphibians. Soil contamination with pesticides should therefore be considered as part of the context of the drastic decline in biodiversity as a whole. ●

GHOSTS FROM THE PAST

Widespread occurrence of pesticides in organically managed agricultural soils, study from 2021



RESIDUES IN KENYA

TOXIC SIDE DISH

Pesticide use leads to residues in food to which many people around the world are exposed. Kenya is no exception. Global trade of pesticide products and produce means that contaminated food is every consumer’s responsibility.

Chemical residues can be problematic to animals and humans. The daily intake of pesticide-contaminated food can pose severe health risks. Sensitive groups such as pregnant women or children are particularly at risk. To protect consumers from pesticide residues in food, governments are usually mandated to take regulatory action for public health protection. Legislation generally provides for the limitation of residue levels that may be allowed in food items entering or leaving various countries. These maximum residue levels (MRLs) are set by authorized regulatory agencies almost everywhere in the world. Since 1963, the United Nations publish the Codex Alimentarius - a collection of standards for food safety and product quality. The maximum residue levels contained therein are considered an important international reference. Nonetheless, there are big differences in the maximum legal intake quantity of pesticide residues depending on the country and region.

For each approved active ingredient, the European Union specifies the maximum concentration of residues to be legally permitted in various food. If goods exceed the limits, they may not be placed on the market. Kenya’s regulatory authority, the Pest Control Products Board (PCPB), in most cases adopts the same MRLs that are used in Europe or the USA, where standards are less strict. This is worrying because local diets are different in Kenya. Maize, a staple food is consumed in greater quantities in Kenya than it is in Europe and should therefore have stricter limits for pesticide residues. Legislation should be strengthened through a local human


health risk assessment taking into account the Kenyan dietary intake and consumption patterns.

Maximum residue levels are based on the cultivation practices, the toxicity of the active ingredient, and food consumption. Baby food must meet stricter specifications. In 2018, under the requirements of the USAID FOODSCAP Project, 1,139 samples of fresh produce intended for export and local markets, were tested by Kenya Plant Health Inspectorate Service (KEPHIS). Pesticides were detected on 46 percent of the samples while 11 percent had residues exceeding EU maximum residue levels. The food items with the highest residues were kales, peas, and capsicum.


Health experts criticize the absence of maximum legal limits for multiple residues in food. A further criticism is that manufacturing companies can circumvent regulations. For example, if active ingredients are at risk of losing their EU approval for health reasons such as being classified as carcinogenic, their maximum residue level for food consumed in the EU is automatically lowered to protect human health. Usually, the limit is lowered to 0.01 milligrams per kilogram, which also applies for imported goods. Pesticide manufacturers who fear a ban of one of their active ingredients for health reasons often just let EU permits expire. This way, they can apply for “import tolerance”: A higher MRL set for imported products to meet the needs of international trade. EU law forbids granting this for pesticides that have lost their approval because of health effects.

The EU has a tighter regulation than many non-EU countries. In Japan, for example, almonds may be contaminated with one milligram glyphosate per kilogram – which is ten times as much as the EU permits. In the Eastern Mediterrane-

The EU has set strict rules for maximum residue limits. However, just like for the approval processes, it fails to take into account the effects of multiple residues



SOMETHING ALWAYS STICKS
Residues of banned pesticides in imported fruits in Switzerland in 2017



an, an area that is home to nearly 680 million people and includes countries from the Middle East to Central Asia, maximum residue levels have been exceeded in up to 61 percent of food samples over the past 15 years. Repeatedly, residues of pesticides that have been banned for a long time are detected there.

Brazil is another example for a lack of efficient regulation that imposes on its population maximum residue levels in food that are two or three times higher than the maximum residue levels in the EU. In 2019, 23 percent of all samples exceeded already high national maximum residue levels according to the official Brazilian residue report. EU-banned active ingredients have also been detected as residues in Brazilian cereals, fruits and vegetables. As export goods, these pesticide residues end up again in Europe or other regions. Put differently: A pesticide which is forbidden in Europe can be exported to a third country, used on crops, and then imported back as a residue on fresh produce.


In Kenya in 2020, a total of 25 different active ingredients were found in tomato and kale samples – 51 percent of the detected active ingredients were already withdrawn from circulation in the EU long ago. Of the total of 25 samples, 60 percent exceeded the EU maximum residue levels. It is alarming in particular, because these two vegetables are often bought from local markets in rural areas and in the city and are used in many Kenyan dishes. However, there is limited information on the actual health risks associated with consumption of such commonly consumed foods.

The increase in pesticide use requires necessary safe-

Many pesticides in Nigeria are proven to cause chronic health effects and show very high environmental toxicity

A European ban on toxic pesticides does not translate into their immediate disappearance. In the last years, a increase to exposure can be observed

guards to control how they are used. Independent, scientific evidence should inform policy decisions and regulation and there should be stronger political efforts to restrict or remove the most toxic pesticides – especially in Kenya and other developing contexts. Nationwide training on the proper use of pesticides is one part of any risk mitigation strategy. However, increasing knowledge and the implementation of integrated pest management approaches – where toxic pesticides are used as a last resort – as well as strong support for non-chemical plant protection, is necessary to reduce pesticide residues in food and enhance food safety. ●



BIODIVERSITY

EXTINCTION IN FULL SWING

Experts have been warning for years that biodiversity is at stake. Pesticides have been identified as one of the causes for why the abundance of animals and plant species is deteriorating so quickly and disastrously.

A significant loss of biodiversity has been observed in the European agricultural landscape for many years. For example, populations of field birds and meadow butterflies have declined by more than 30 percent since 1990. The structure of agricultural landscape is the most common cause, mainly the size of fields, lack of landscape features such as hedgerows or ponds – and the usage of chemicals such as artificial fertilizers and synthetic chemical pesticides.


There is a consensus that pesticides play a significant role in biodiversity loss – they harm biodiversity directly and indirectly. The control of weeds by broad-spectrum herbicides such as glyphosate leads to a decimation of flowers and blossoms and thus to a shortage of food for insects that feed on flowers and wild herbs. In 2017, the total sales of glyphosate are estimated at more than 46,000 tonnes across the EU. In the same year, glyphosate sales were highest in France followed by Poland and Germany. In the latter Country 40 percent of all agricultural land is treated with the herbicide.

2021 study results show the impact of pesticide applications on biodiversity. A German institute recorded and analyzed floral diversity relative to methods of cultivation in the agricultural landscape. The ratio in terms of species diversity and coverage in fields and of the actually flowering species and their flowering intensity was 3 to 52 to 100 from fields managed conventionally for many years to fields managed organically for many years and fields which never faced the usage of chemical pesticides. Because wild plant species in fields are important sources of nectar and pollen their decline as a result of intensive management with herbicides can also be expected to have a significant impact on the diversity and abundance of insects in the arable-dominated agricultural landscape.

The sharp decline in insects in agricultural landscapes has been documented by many studies. The population of grassland butterflies in European countries has decreased by about one third between 1990 and 2015. EU Red Lists show that almost 10 percent of bees are threatened with extinction in Europe mainly because of agricultural practices including the use of pesticides and fertilizers. The most widely used insecticides are neonicotinoids, which are very toxic to insect pollinators like bees. Therefore, 4 out of 5 active ingredients are now only allowed with exceptional approval. Bees and other pollinators can be exposed to pesticide through different ways. For example, pollen and nectar from pesticide treated plants may contain residues: A study published in 2017 found pesticides in honey from across the world. 75 Percent of all honey samples contained at least one neonicotinoid. More than one third of honey samples were contaminated with concentrations of neonicotinoids like imidacloprid that are known to be detrimental to bees. Similar substances were detected in a study the German environmental organization BUND conducted. More than half of the samples – ordinary honey sold in German supermarkets – were showing residues of pesticides like acetamiprid or thiacloprid. Based on the available data, thiacloprid has been classified as likely to be carcinogenic in humans. Studies found that a chronic exposure to thiacloprid significantly impaired honeybees’ foraging behavior, immune system and navigation – or kills them directly.

There is a growing body of research showing pesticides can become more harmful when mixed – even when components were combined at concentrations below its individual no-observed-effect-concentration (NOEC). For example, some fungicides can increase the toxicity of pyrethroid insecticides for bees. Scientific knowledge of pesticides suggests that it is insufficient to reduce the amount of pesticides used – even in very small quantities many substances can endanger biodiversity. It is more decisive how toxic the active


Soils contain nearly a quarter of the planet’s diversity. Pesticides often harm organisms that are essential for their conservation




THE WORLD IS LOSING ITS DIVERSITY

Environmental impact of pesticide use


Floral diversity and flowering intensity in relation to cultivation methods on arable land



Main drivers of insect decline



In sharp decline: Seasonal distribution of insect biomass (grams per day), representative of Western European low-altitude nature protection areas embedded in a human-dominated landscape



ingredients are for certain animals and plants. A study of the University of Landau (Germany) found that the total amount of insecticides used in the USA was reduced by 40 percent between 1992 and 2016. Fish, mammals and birds benefited from this, as this decrease was mainly due to decreasing use of certain classes of insecticides such as organophosphates and carbamates, which are problematic for these groups. However, a different picture emerges for invertebrates such as crustaceans or insects and especially pollinators insects such as bees. Despite the decline in the amount of insecticides, toxicity for these groups more than doubled between 2005 and 2015.


Factors like the amount applied per acre or other unit and the persistence of pesticide residues in water or soils shed light on how certain pesticides cause adverse effects on nature. In addition, efficacy should not be underestimated: Highly effective pesticides can have the same hazard potential as older substances in higher doses. For this reason, European civil society organizations are not only calling for a reduction in the amount used, but also for a ban on particularly harmful pesticides.

The EU did not meet its latest target to improve the situation of protected species. More than two-thirds of species assessments result in a concerning conservation status

The use of organic management practices in field cropping has big effects on floral biodiversity: on longstanding organic farms it is 17 times higher compared to conventional fields

UNPRECEDENTED DECLINE

Conservation status of species, by member state level, in percent



BENEFICIAL INSECTS

SMALL CREATURES WITH BIG IMPACTS

Beneficial insects provide enormous ecosystem services to Africa. However, to continue benefitting from these natural processes, sustainable farming and environmental management practices need to be promoted.

Africa is among the most biodiverse regions in the world. It is home to more than 25 percent of the global biodiversity and has eight of the 25 biodiversity hotspots. Regions such as the Congo-basin, South Africa and Madagascar are considered megadiverse. Ecosystem services from this rich biodiversity have enormous potential to contribute to better livelihoods. There are nearly 7 million different insect species on the continent that provide pollination, pest control, decomposition, and maintenance of overall biodiversity.

Generally, insects are viewed as pests. However, only a small subset of insects are either pests of crops or carry human and livestock diseases in Africa. Pest populations are kept under check by natural factors, such as parasitoids, predators, and microbes. For instance, a small native micro


wasp, *Cotesia icipe*, can specifically target armyworms and parasitize over 120 invasive fall armyworm larvae and kill them. Other natural enemies like dragonflies, damsel flies, predatory bugs, maggots of hoverflies, or rove beetles aid to keep in check a wide variety of insect pests. However, unsustainable, and intensive farming practices such as misuse of pesticides, monocultures and land use changes can significantly disrupt this natural balance. In addition, invasion by pests which are not native to Africa and changing climates, results in catastrophic outbreaks of crop pests affecting food security, human, animal and environmental health.

Conservation of natural enemies through sustainable farming practices can offer effective management of insect pests with positive economic and health impacts. For instance, release and conservation of a small parasitic wasp, *Anagyrus lopezi*, resulted in effective control of invasive cassava mealybug, *Phenacoccus manihoti*, in over 27 sub-Saharan Africa countries with estimate benefits of over 9.4 billion US dollars. In East Africa, release and conservation of parasitoids, *Diadegma semiclausum* and *Cotesia vestalis*, resulted in successful management of *Plutella xylostella*, a pesticide resistant pest of crucifers with significant economic impacts. Similarly, invasive fruit flies, *Bactrocera dorsalis*, is biologically controlled with parasitoids, *Fopius arisanus* and *Diachasmimorpha tryoni* in Africa. To ensure that countries continue to benefit from these biological control efforts, sustainable and compatible farming practices need to be promoted and scaled.

Beneficial insects do not only protect crops from pests, but they also aid in pollination. Several crops grown in Africa, such as avocado, cucurbits, African nightshades, cocoa, spider plants among others depend on insect pollinators to improve crop yields and quality. Beyond crops, pollinators contribute to maintaining plant biodiversity in natural vegetations. When we think of insect pollinators, most of us are reminded of honeybees, *Apis mellifera*, a major pollinator in the ecosystem. However, there are over 2,600 other species of bees in Africa that contribute to pollination. Some of them, such as stingless bees, also offer diverse and unique honey and honey-related products, like bee wax, to improve income and livelihood of farmers. Other insects such as moths, wasps, flies, beetles and ants also contribute to pollination. Close interaction of pollinators to cultivated crops, makes them vulnerable to unsustainable crop management practices such as pesticide misuse.


Organic fertilizers are a key resource in sustainable ag-

Biological pest control uses other organisms to control pests. It is a safe, species specific and long-term action against target pests and a sustainable way to increase food production in Africa



GRILLED, BOILED OR STEAMED

Number of insect and spider species recorded as being consumed by humans, 2017



riculture, the availability of which is scarce in Africa. Several insects such as house flies, blow flies, soldier flies, dung beetles and flower chafer beetles, thrive on organic waste and recycle them to high value biofertilizer for improved soil health and crop yields. In addition, the insect biomass generated also presents a rich source of protein, fats, minerals, vitamins, well-balance amino acids and other essential elements, which can be used as high-quality ingredients in animal feeds to substitute or replace the expensive and scarce fish and soya bean meals. These insects are an excellent source of oil with their potential use as biodiesel, additives in soap and cosmetic production as well as immunity enhancers.


The emergence of starch rich crops and domesticated livestock and birds has eroded the cultural acceptance of insects as a food option globally. However, in Africa, insects continue to be a traditional food with over 500 insect species consumed. Termites, locusts, long-horned grasshoppers, ants, saturniid caterpillars, crickets, among others, contribute to much-needed proteins, fats and essential minerals. Currently these insects are accessed from the wild during seasonal outbreaks which is increasingly affected by climate change. Wild harvested insects risk contamination from microbes, heavy metals, and pesticides. Recent research efforts in Africa led by the International Centre of Insect Physiology and Ecology (*icipe*) are focussed on science-based mainstreaming of edible insects, through sustainable mass rearing for their year-round availability. The focus is also to

In Africa, insects continue to be a traditional food with over 500 insect species consumed. Termites, locusts, long-horned grasshoppers, ants, crickets, among others, contribute to much-needed proteins, fats and essential minerals

improve the nutrient profiles of edible insects for health and nutrition.

The biosafety of edible insects harvested from the wild and those mass-reared has been led to the development of enabling policies on edible insects in Kenya, Uganda, Rwanda and Tanzania. The positive research outputs and favourable policy environment has attracted private sector investments and youth business development and employment in the edible insect sector in East Africa. ●

Ladybugs eat aphids, but pesticides kill these beneficial insects – agrochemicals contribute to the comeback of harmful insects. Pesticides kill these beneficial insects




WATER

GO WITH THE FLOW

Studies regularly reveal how pesticides contaminate rivers, lakes, coastal waters and groundwater. The pollutants often originate in agriculture and enter surface water via infiltration, surface runoff and drift.

Mixtures of chemicals such as pesticides, biocides, pharmaceuticals and industrial chemicals have been detected in rivers, lakes, and other surface water all over Europe. These pollutions affect the living conditions of aquatic organisms and the general ecological status of water bodies in Europe. Stressors such as climate and land use change or water scarcity make the situation worse. In other parts of the world such as China or South Africa the water quality of rivers, lakes, and groundwater is even more threatened by pesticides. There, pollution has particularly far-reaching consequences because there is less overall availability of freshwater and the water bodies harbour a great deal of biodiversity.



With regards to water pollution by pesticides, one figure in the European approval process is particularly meaningful: the regulatory acceptable concentration (RAC) per active ingredient. The assumption is that the harmful effects of pesticides on aquatic life are low as long as this concentration is not exceeded in the water. Despite this requirement, small streams, which make up a large proportion of European watercourses, are regularly contaminated with pesticides, according to environmental monitoring. They are often located in the middle of agricultural land and thus particularly exposed to pesticides, as a recently published study by the Helmholtz-Centre for Environmental Research together with the German Environment Agency proved.

A study by the European Environmental Agency shows that levels of pesticides exceeding national thresholds were measured in up to one-third of all reported monitoring sites in European surface waters from 2013 to 2019. The pesticides that most often exceed thresholds are the insecticides imidacloprid and malathion, and the herbicides metolachlor and metazachlor.


At global scale, the situation is even more alarming. A study from scientists from a German University provides a comprehensive meta-analysis of 838 peer-reviewed studies that evaluates the exposure of surface waters to insecticides. Among the 11,300 insecticide concentrations detected, more than half exceeded their threshold levels – so the biological integrity of global water resources is under substantial threat. Because residue analyses are too expensive for many local scientific institutes and there is a lack of national monitoring data in the Global South, one can assume that the figures would likely be even greater with more data. What is already clear, though, is that global chemical pollution levels have exceeded planetary boundaries.

It is certain that agricultural pesticides are a crucial environmental stressor for insects in small water bodies. Studies show that in polluted streams in Germany, populations of sensitive species such as dragonflies and caddisflies decrease significantly. But not only small streams are at risk: Pesticides ultimately end up in the sea via rivers. Environmental experts have been studying the presence of pesticides in harbor seals and other marine mammals in the EU LIFE APEX project. The results show that the pesticides that are particularly problematic are those that persist for long periods of time in the environment and can accumulate and pass from one species to the next through the food chain. One example is hexachlorobenzene (HCB). This pesticide, which was originally used as fungicide, has been banned in European agri-

Many factors contribute to the bad ecological status of water bodies. Exceeded legal limits on pesticides concentration are one of them. In Germany's small water bodies for example 81 percent of all monitoring stations showed levels above the limit

INFILTRATED ENVIRONMENT

Chemical status of all surface water bodies, selected EU Member States



culture for 40 years. And yet dolphins, porpoises and seals in European seas are still heavily contaminated today.

Through rain infiltration or leaching, pesticides move into the ground water, where they degrade slowly. A study in Germany detected active ingredients at almost one third of the monitoring sites. Degradation products were found at even 58 percent of the monitoring sites. In Italy, about one third of the groundwater bodies investigated showed pollution by pesticides. Frequently detected pesticide compounds in surface water and groundwater include glyphosate and its degradation product AMPA. A mandatory threshold also for degradation products would allow better regulation – so far there are only non-binding recommendations. Another important measure that could protect water bodies from pesticides is establishing continuous riparian buffer zones, which additionally provide an important habitat for plants and migration corridors for animals. Such riparian buffers in which the use of pesticides is prohibited are mandatory only in a few countries. In many regions of the Global South they are practically not feasible at all, as the agricultural area is often smaller than the required width of the riparian buffer zone.


Environmental experts point to the need of an agricultural turnaround: A comprehensive reduction of pesticide pollution of water can only succeed through restructuring conventional agriculture towards less use of chemical pesticides. Protecting soils and improving their quality could prevent erosion which in turn reduces the runoff of pesticides. ●

Active ingredients banned due to their hazardous properties stay a long-term problem – even long after their ban

According to the European Environment Agency, many lakes, streams, transitional and coastal waters are not in good ecological status. And even groundwater is polluted

LONG LASTING IMPACTS ON MARINE INHABITANTS

Contamination of mammals with the fungicide Hexachlorobenzene (HCB) banned in 1981, in nanogram per gram wet weight



LONG-RANGE TRANSPORT

GONE WITH THE WIND

Pesticides rarely stay in the place where they have been applied. Wind can move dust, particles, and droplets to residential areas close to agricultural land – or carry it to places many kilometres away. Approval processes are largely ignoring this problem.

When pesticides are applied with spray nozzles, droplets or mist can be blown by the wind onto neighboring land. This phenomenon is called pesticide drift. Incorrectly adjusted and inappropriate nozzles or excessive speed of the spray vehicle intensify the effect. Active ingredients may also travel much longer distances, from a few hundred metres to over 1,000 kilometres. This is called “long-range transport”. Active ingredients can rise into the air; because of ground warming, evaporation or adhering to tiny dust particles being blown up by the wind from uppermost soil layers. In this case, air currents distribute small suspended particles – so-called aerosols – in all directions. Cooling and rain cause them to sink back to the ground. They can end up almost everywhere: in nature reserves, in city parks and in human lungs.


The possibility of long-distance transport of pesticides has long been known. As early as 1999, a study collection drew attention to the fact that 30 pesticides were found throughout Europe, in some cases at measuring points far away from where they were applied. For a study published in 2020, two German NGOs (Bündnis für eine enkeltaugliche Landwirtschaft and Umweltinstitut München) examined pesticide contamination of air. At 163 sites throughout Germany – including protected

areas, cities and organic fields – traces of 138 pesticides were detected.

30 percent of the substances found are not or no longer permitted in Germany, for example DDT, a long-lived organic compound that is difficult to degrade and prohibited in most western countries since decades. Cocktails of 5 up to 34 pesticides and their degradants were found at three quarters of the sites. Glyphosate, the most widely used herbicide in the world, was detected at all sites that were equipped with technical filters. This is significant because it disproves the assumption that glyphosate does not spread through the air – glyphosate and all its salts are considered non-volatile, which is why the European Food Safety Authority (EFSA) has so far ruled out the possibility of long-range glyphosate transport.

Another 2020 study examined airborne pesticide concentrations at 50 sites across France over a 12-months period. Glyphosate was detected at 80 percent of the sites investigated. This is further evidence for large distance transport of glyphosate through the air. The fact that long-range transport and drift occur worldwide is demonstrated by other recent studies. To assess possible contamination of non-target areas in South Tyrol, 71 grass samples of public playgrounds and schoolyards located next to intensively managed apple and wine orchards were examined. At least one pesticide and sometimes even pesticide cocktails were detected in 96 percent of the samples. The majority of the

Residue data were analyzed from grass samples by an international research group. They found endocrine active substances – some of them are suspected human carcinogens



TOXIC LONG-HAUL FLIGHTS

Monitoring stations in Germany for airborne pesticide mixtures, and distance of detected residues to their presumed application area

- between 100 and 1,000 metres
- more than 1,000 metres

□ Nature reserves

Brocken, Harz National Park:
12 pesticides transported over a large distance found, partially in considerable quantity

Bavarian forest:
5 pesticides transported over a large distance found, including glyphosate and the recently banned chlorothalonil and chlorpropham

Study from 2020

detected pesticides are classified as endocrine disruptors, which can affect the health of humans and animals, even in minuscule amounts. Another example from the USA shows air pollution probably caused by pesticide drift. According to a 2021 study, more than one million acres of soybeans and at least 160,000 acres of a conservation area were affected by exposure to the herbicide dicamba from adjacent agricultural fields.

For years, civil society organizations in South Africa and other countries have been advocating for mandatory buffer zones as a risk mitigation measure. A new measure was also imposed in France to protect residential areas from drift of hazardous pesticides – farmers must respect now a buffer zone of 20 metres.

A national air monitoring program of pesticides exists only in Sweden. And in approval processes for pesticides and active ingredients, little attention is paid to the phenomenon. The risk of a possible long-range transport is only estimated theoretically. A verification of the contamination in practice, however, does not take place.

The estimated amount of pesticides that people can consume on a daily basis without any immediate risk to health is only based on digestive tract absorption and only for a single active ingredient at a time. In contrast, pesticide exposure through drift and long range transport takes place primarily through the respiratory tract – and the long term effects of pesticide cocktails entering the human body through the lungs are still largely unknown. ●

Organic farms that forgo pesticides are under threat from volatile substances – wind carries them onto organic fields and this can threaten their business

Air quality under threat: A cocktail of five to 34 pesticides was found at 75 percent of all monitoring stations

ORGANIC FARMING UNDER PRESSURE

Drift and long-range transport of pesticide

250,000 EUR is the cost each year for a large medium-sized organic business to check whether their products have been contaminated

30 percent of active ingredients detected in long-range transport analyses between 2014 and 2019 are no longer permitted at the time of measurement, including the insecticide DDT, which has been banned for decades

INSECT DECLINE

AN ECOLOGICAL ARMAGEDDON

Insects provide pollination services to flowering plants, control pests and ensure abundant harvests. For a long time their populations have declined dramatically – causing disaster for humans and nature. Pesticides are considered a major reason for the decline.

Insect populations have declined sharply in recent decades. These downturns are of direct concern to humankind as we rely upon insects to deliver vital ‘ecosystem services’ such as pollination, recycling of nutrients and pest control. A review by the University of Sydney in 2018 compiled information from research studies in various regions. It found that the populations of 41 percent of species are in decline, and one-third of all insect species are threatened by extinction. While cautioning that the available evidence was relatively thin, the researchers estimated that total insect biomass is declining by 2.5 percent a year. Most of the research studies they included in their review came from Europe, some from North America and only a few from Asia, Africa or Latin America. Some examples: UK butterfly populations have fallen by about 50 percent since 1976, the biomass of flying insects in German nature reserves declined by 76 percent in the 27 years to 2016. In North America, populations of the Eastern monarch butterfly have fallen by 80 percent in 30 years, and in the Netherlands numbers of caddis flies fell by 60 percent between 2006 and 2016. There are many data gaps, particularly for tropical regions, but the evidence suggests that insect declines are a global phenomenon, and that they are ongoing.


There is broad agreement amongst scientists that insect declines are driven by a range of factors, including habitat destruction, climate crisis, light pollution, increasing fer-

tilizer use, and the impacts of invasive species. Pesticides play a key role as well. Impacts of pesticides on insect populations have been examined in most detail for butterflies, a group of insects for which exists relatively good population data. For example, organic farms have been found to have more butterflies than non-organic neighbors, and pesticide-treated gardens had about half as many butterfly species as untreated ones. Use of neonicotinoid insecticides in particular have been found to correlate with patterns of butterfly decline, in both UK and California. However, it is not possible to accurately specify to which extent the decline is linked to the use of pesticides, not least because habitat loss, farming intensification and pesticide use are all strongly correlated with another.

The impacts of pesticides on the environment were first highlighted in 1962 by Rachel Carson in her book Silent Spring, which drew attention to the problems being caused by the extensive use of early insecticides such as DDT (dichlorodiphenyltrichloroethane) and organophosphates. Although these early chemicals were eventually banned in most countries, they have been replaced with successive generations of new compounds, many of them much more toxic to insects. For example the neonicotinoid insecticides, introduced in the 1990s and now the most popular insecticides in use globally, are approximately 7,000 times more toxic to insects than DDT.


According to their effect different pesticides have a different impact on insects: Even though insecticides should protect plants from pests they harm all insects, both the pests and beneficial insects. Since pesticide applications

Neonicotinoids like Imidachloprid have been described as a worldwide threat to biodiversity – but some EU countries continue to use and export them



SELF-HARMING BEHAVIOR

Dependence of agricultural production on pollination by animals, 2012*



kill natural enemies of crop pests (insects such as ladybirds, hover flies and lacewings), populations of crop pests such as aphids often bounce back rapidly.

But also fungicides and herbicides are harmful to insects. For example some fungicides act synergistically with insecticides, rendering them more toxic if an insect is exposed to both at the same time. The herbicide glyphosate has recently been found to be harmful to bees, damaging their beneficial gut microbes and also affecting their learning abilities. Further, herbicides remove weeds such as wildflowers and food-plants which removes vital resources for insects for their larvae, thus indirectly impacting insect populations.

Systemic insecticides such as neonicotinoids contaminate soils and are taken up by the roots of wildflowers, so contaminating the nectar and pollen. Neonicotinoid insecticides have a range of sublethal impacts on bees, including impaired learning which interferes with communication and navigation; reduced immune function rendering them more susceptible to diseases; and reduced fecundity. A recent study found neonicotinoid insecticides in 75 percent of honey samples collected from around the world. Honey samples often contain not just neonicotinoids but a cocktail of ten or more pesticides, often including other insecticides, herbicides and fungicides. If honeybees are being exposed to these mixtures then it is very likely that thousands of other species of beneficial pollinating insects are also consuming them when they visit flowers. All these impacts are not taken into account enough in the regulatory process. Some negative impacts on pollinators are not even detected by regulatory studies. A progressive decline in insects threatens


Pesticides pose a threat to insects and to the economy: The pollination services provided by insects are worth 153 billion Euros per year

In absolute terms, the losses seem relatively limited. Many cereals are not dependent on pollination – unlike the majority of fruit and vegetable species from which we obtain vital vitamins and minerals


vital ecosystem services such as pollination, recycling, and biocontrol of pests, as well as removing a vital component of food webs, and ultimately endangers human wellbeing through the quality and quantity of our harvests. ●

HELPERS NEED HELP

Threatened crop loss in absence of pollination by animals



Global decline in insect species, 2009–2019



HIGHLY HAZARDOUS PESTICIDES

A GLOBAL HUMAN RIGHTS CONCERN

Substances that are proven to present a particularly high level of acute or chronic risk to health or the environment are commonly referred to as **Highly Hazardous Pesticides (HHPs)**. Far too rarely are these substances withdrawn from circulation – especially in the Global South they cause great harm.


To identify HHPs, the Food and Agriculture Organization of the United Nations (FAO) and the World Health Organization (WHO) have outlined eight criteria: Pesticides are considered to be highly hazardous if they have an acute lethal effect, cause cancer or genetic defects, impair fertility, or harm unborn children. Likewise pesticides are classified as highly hazardous if they cause serious or irreversible damage to health or the environment under normal conditions of use or are listed in internationally binding conventions like the Stockholm Convention on Persistent Organic Pollutants, the Rotterdam Convention, or the Montreal protocol.

Although the FAO and WHO developed these criteria, they have not published an official list that includes all HHPs used worldwide yet. This makes it challenging for governments, agricultural extension agents, distributors, and appliers to identify and replace HHPs with less hazardous alternatives. The international Pesticide Action Network (PAN) has filled this gap and has published a periodically updated HHP list since 2009. It takes into account environmental criteria as well as additional human health impacts compared to WHO and FAO.


For years, studies have shown that HHPs cause great damage especially in countries in the Global South, and yet massive amounts of these specifically harmful pesticides are still applied to a vast extent there. In 2018, 40 percent of all pesticides used in Mali were highly hazardous, in Kenya 43 percent at the same time. In 2021, even 65 percent of all pesticides used in four states of Nigeria were highly hazardous. In Chile, one quarter of all 400 active ingredients registered were HHPs in 2019, and in Argentina as many as 126 out of a total of 433. The use of HHPs in agriculture is also widespread in Eastern Europe, the Caucasus and Central Asia. Investigations could show that between 2019 and 2021 more than 70 HHPs were used in Georgia, Kyrgyzstan and Ukraine, and as many as 95 in Armenia. Even though the EU has banned many HHPs, some specifically dangerous pesticides remain in use, even though they should be substituted according to EU regulations.

In many countries, the system of pesticide regulation is inadequate. Capacity with regards to quality and use control, advisory services and monitoring of pesticides are often insufficient or even entirely lacking. Many of the workers applying the pesticides are also poorly trained or not trained at all: The lack of safety trainings frequently leaves them unaware of the health hazards involved in handling pesticides. A lack of information about hazardous substances and difficulties in accessing disposal centers for empty pesticide containers impedes the return process. In some

Contaminated food, a large number of highly hazardous substances and hardly any means of protection: NGOs call such a situation a humanitarian catastrophe



PUTTING THE WORLD AT RISK
Percentage of pesticides considered as highly hazardous, by country



countries, disposal centers do not even exist. And in many cases there is not even access to personal protective equipment or hot climate makes wearing such impossible which creates additional problems. This results in a high number of injuries and deaths: 95 percent of 385 million people who suffer from unintended pesticide poisoning each year live in the countries of the Global South. United Nations experts have considered HHPs a global human rights concern for a long time: Pesticides threaten among others the right to live in dignity, the right to bodily integrity, and the right to a healthy environment. Also, pesticides are often applied disregarding mitigation measures like buffer zones to protect surface waters, or specific spraying times to protect pollinators, and even though these measures are practically not feasible in many regions, the pesticides still remain on the market.


Despite their dangers, using HHPs seems normal these days – but it does not have to be. Many regional projects in both the South and the North have demonstrated that agroecological farming practices are a viable alternative. However, this transformation can only succeed if governments and the international community set appropriate priorities. It is particularly important to raise awareness of the risks of pesticides and to push for the development of non-chemical alternatives. Key elements include research funding, and the collection and dissemination of information on viable alternatives to HHPs, ranging from ecological and cultural management measures to biological

There are criteria for identifying HHPs, but no international agreed convention or protocol addresses all of them. Not even 4 percent of all pesticides used globally are regulated by binding international conventions

Regulatory measures often correlate with the country's prosperity. Civil society organizations are calling for a global legally binding mechanism for the lifecycle management of pesticides

cal control measures and as a last resort a restrictive use of biopesticides.

A progressive ban on HHPs was recommended by the FAO as early as 2006. Developing safer alternatives is the goal of the Strategic Approach to International Chemicals Management (SAICM), which aims to reduce the usage of Highly Hazardous Pesticides. Nevertheless, there is still no globally binding legal framework that addresses pesticides in their full scope – from production to use to disposal, and with strict deadlines for phasing out HHPs. ●



GENETIC ENGINEERING

MODIFIED CROPS, MORE PESTICIDES


Genetically modified crops were supposed to reduce the use of chemicals in agriculture, reduce workloads, and increase crop yields. These promises could not be kept.

More often than any other substance, glyphosate has been at the center of many controversial debates about pesticides in recent years. In 2017, EU Member States had voted to extend the license for the herbicide for at least five years, despite cautionary voices and demonstrations in numerous countries. How does the herbicide actually work? In short: Glyphosate is applied to food and nonfood field crops such as soybeans and field maize. Glyphosate inhibits the EPSPS enzyme, which is required in plants for the production of vital amino acids. This interrupts the metabolism – and the plant dies. Genetically modified crops are protected against this interruption of the metabolism and can therefore continue to produce amino acids and survive despite sprays. For this reason a genetically modified soybean in its growth phase can be

treated with glyphosate without being harmed – while all surrounding plants, that compete with it for water, space and nutrients, die. In times before genetic modification, competing plants usually had to be controlled either by pre-emergence herbicide application, by crop rotation or manual weeding.

Today, 74 percent of soybeans grown worldwide are genetically modified. The increased use of genetically modified organisms (GMO) has been associated with a massive increase in glyphosate use. From 1995 to 2014 the agricultural use of glyphosate in the US rose ninefold, reaching 113,000 tonnes per year – one-third of the total amount of herbicides applied. From 2012 to 2016 an average of approximately 127,000 tonnes of glyphosate were applied to 120 million hectares annually. Most glyphosate was applied to soybeans (53,000 tonnes), maize (43,000 tonnes) and cotton (9,000 tonnes). Globally, the total use of

Investments and buyouts:
A few companies have taken over the market
for seeds, especially in the Global South



glyphosate rose almost 15-fold, from 51,000 tonnes in 1995 to 747,000 tonnes in 2014. This increase correlates with the expanded cultivation of GM soy in Latin America. After its introduction in Argentina in 1996, the glyphosate volume there had doubled within just one decade. In Brazil, herbicide use in soybean cultivation tripled from 2002 to 2012 to 230,000 tonnes per year, mainly due to glyphosate. Despite the drastic increase in herbicide rates applied, yields per hectare increased by only about 10 percent. Brazil and Argentina are now among the countries with the highest herbicide consumption in the world, in third and fourth place globally after China and the USA.

Intensive use of glyphosate has led to the appearance of glyphosate-resistant weed species worldwide. First reports from Delaware, USA, made global headlines in the year 2000. They found that the Canadian horseweed could no longer be controlled with glyphosate. By 2012, herbicide resistant weeds have already spread across 25 million hectares of arable land in the United States. There are now 53 weed species that have developed glyphosate resistance, including amaranths in cotton and soybean crops. In order to combat such weeds less sensitive to glyphosate, farmers have increased glyphosate application rates and the use of other herbicides was intensified again as well.


Another genetic modification intended to contribute to pesticide reduction was the insertion of specific DNA sequences into crop plants to enhance their resistance to insect pests: A gene transfer from the bacterium *Bacillus thuringiensis* leads to the formation of proteins known as Bt toxins in the plants. Those proteins are lethal to several types of insects. Insect-resistant crops were cultivated in the mid-1990s for the first time, nowadays they make up 57 percent of all genetically modified crops grown around the globe, predominantly maize and cotton. The fact that plant-incorporated toxins in all parts of the plant act as insecticides throughout the entire vegetation period has consequences for the environment. For example, butterflies and other insects can be harmed. And just like the weeds in soybean cultivation, pests also develop resistance.

In the USA, specimens of the Western corn rootworm are already resistant to more than one Bt toxin. At the beginning of Bt crop cultivation, the number of pesticides used actually decreased. But only impermanently: Sales of insecticides in corn production in the US have increased significantly. In 2018, Indian farmers spent 37 percent more money per hectare on insecticides than before the introduction of genetically modified cotton in 2002. In addition, the cost of seed and fertilizer increased.

These complaints are not new: Already more than ten years ago, twenty civil society organizations from India, South Africa and all over the world stated in their declaration “A Global Citizens Report on the State of GMOs” that genetic engineering has failed to increase food crop yields but has vastly increased herbicide use and the growth of resistant weeds. While big companies gaining seed market control and pushing up prices, farmers have to go into debt. The high levels of indebtedness among farmers is, for example, thought to be behind many of the hundreds of thousands deaths by suicide of Indian farmers over the past years.

PATH COMPANION
Cultivation area of genetically modified soybean (GM soy) in North America and South America in 2019, in hectare

- Number of herbicide-resistant weeds
- Share of GM soy in total soybean cultivation
- Increase in herbicide use between 2009 and 2019
- Cultivation area of GM soybean



*Based on official figures and UN estimates

© PESTICIDE ATLAS 2022 / FAOSTAT, ISAAA, WEEDSCIENCE

There has been a substantial increase in the usage of herbicides in soybean production. It appears to be linked to genetically modified beans

SMALLHOLDER FARMERS

NEW MARKETS, LESS REGULATION

In Africa, fewer pesticides are used than in other regions of the world. Nevertheless the 33 million smallholders are increasingly becoming the focus of pesticide companies. There they also sell what has been banned in the European Union.

In 2015, the African agrochemical market was valued at about 2.1 billion US dollars. It accounts for only 2 to 4 percent of the global usage. According to the Food and Agriculture Organization of the United Nations (FAO), an average of 0.4 kilograms of pesticides were used per hectare of cultivated land in Africa in 2019. This is less than the 3.7 kilograms in North and South America. But the African market for pesticides is projected to witness high annual growth rates, for example in West Africa. Pesticide use increased there by 177 percent between 2005 and 2015. In the same period total pesticide imports into the region roughly tripled, with particularly rapid growth in the three largest agricultural markets – Ivory Coast, Ghana, and Nigeria. Coupled with population growth, and the need to improve productivity, pesticide companies are increasingly seeing the 33 million small farmers on the continent as an attractive market.

Major players in the African pesticide market are Adama Agricultural Solutions, Sumitomo Chemicals, UPL Limited, and Bayer AgroScience AG. Companies use specific selling strategies to unleash market potentials in African countries. In Kenya, for example, social media, local radio stations, and broadcasts in local dialects are some of the most used mediums for product advertising. The documentary film “The Food Challenge” shows that prior to the COVID-19 pandemic,



dominant pesticide companies frequently sponsored agriculture trade shows.

Depending on the crop, capital availability, and geographic location, farmers use pesticides very differently. Field studies from Mozambique and Zambia show the widespread use of Highly Hazardous Pesticides (HHPs) – according to a Michigan State University study, 76 percent of farmers in Zambia and 87 percent in Mozambique use them.

Small scale farmers and farm workers are particularly vulnerable when it comes to pesticide use. Mitigation measures are not practical because they are expensive or the farming context does not make risk management possible. In regions such as Africa, Asia and Latin America, smallholder farmers cannot afford proper backpack sprayers, masks, protective clothing, and gloves. In addition, buffer zones are not maintained because farm sizes are small and closely situated to each other and other homesteads. Pre-harvest intervals are often not known by the farmers or ignored because there is financial pressure to sell produce. Pesticides are also decanted from one container to another after they are bought from the agro-vet store, which means that instructions on how to use a product ‘safely’ have been removed. Civil society organizations blame weak regulations and the lack of information by industry for exposing farmers to these risks.

Further, different scientific studies show that pesticide markets in various African countries are not regulated in a

Small-scale farmers are highly exposed to toxic pesticides. They are not able to afford full protective gear or follow label instructions despite receiving information from the Ministry of Agriculture and pesticides sales agents



way which protects farmers’ health and the environment. Another problem is that rules, laws, approvals, and controls could not keep pace with the increasing demand for pesticides – that is why a lucrative market for cheap generic and illegal pesticides has developed. Industry and academic sources estimate that up to 20 percent of the African market, and as much as 34 percent the West African market, are illegally produced and traded. In extreme situations, that number exceeds 40 percent of pesticides. Empty packaging and canisters are also filled with counterfeit products and sold as originals – with serious risks for farmers and the environment.


Civil society organizations are demanding stricter rules for pesticide market approval and authorisation informed by local data. They want governments to explore options to make regulatory risk data more transparent and accessible. Pesticide sales should be regulated and monitored accordingly, by independent authorities. Qualification criteria for agrovet sellers should be established and implemented.

Plant pathogens and pests are a major threat to the African farming sector, the incomes of producers and ultimately, achieving of the human right to food. Smart answers are needed to balance crop protection, which is necessary to ensure sufficient harvest, with human and environmental health: For example, investments in agroecological strategies and evidence-based knowledge sharing amongst farmers, experts, scientists, and policy makers. In some parts of the world this is already taking place. As a first step, organic

NGOs criticize a lack of safety standards in low-income countries. In Uganda every fourth shop sells repackaged pesticides

Five in every six farms in the world consist of less than two hectares – which produce roughly 35 percent of the world’s food. In most cases the farmers suffer from poverty

farming has gained popularity for years. The organic acreage in the Middle East and in Africa is increasing as well. But these are only small steps on a long way. Even though scientists in the last years strongly point to the potentials of agroecological and organic farming methods these are still hardly supported by African governments. ●



IMPORTS AND EXPORTS

BANNED BUT SOLD ANYWAY

Many pesticides are banned in the European Union. It is illegal to use them in EU Member States, yet it is allowed to produce and export them to third countries – where they pose great risks to people and their environment.

According to market forecasts, the number of pesticide exports to countries in the Southern Hemisphere will continue to grow. The five largest pesticide companies –including Bayer, BASF, and Syngenta – already generate more than one-third of their pesticide sales from active ingredients classified by the Pesticide Action Network (PAN) as highly hazardous. According to the World Health Organization (WHO) and the Food and Agriculture Organization of the United Nations (FAO), Highly Hazardous Pesticides (HHPs) present particularly high levels of acute or chronic hazards to humans and the environment. For this reason, many of these pesticides are no longer authorized in the European Union.

However, European companies are still allowed to sell these pesticides – namely to countries outside the EU. This


practice creates double standards. In 2018 and 2019, EU countries and the United Kingdom approved the export of a total of 140,908 tonnes of pesticides that are banned from being applied in European fields because of unacceptable health and environmental risks. Furthermore, European corporations like the German companies Bayer and BASF sell pesticide products locally in third countries with active ingredients banned in the EU. In South Africa and Brazil, they sold products containing at least 28 such active ingredients, according to a 2020 study. Some of the hazardous pesticides exported from Europe find their way back as residues in imported food. Residues of 74 pesticides banned in the EU were found in food tested on the European market in 2018 – 22 of which were exported from Europe that same year.

Brazil today is one of the largest consumers of pesticides in the world and imports most of the pesticide active ingredients from abroad, including from EU countries. In 2019 these included at least 14 highly hazardous active ingredients no longer approved in the EU. Among them were BASF's fipronil, which is highly toxic to bees, nerve damaging chlorpyrifos from Portugal's Ascenza Agro SA, as well as Germany's Alzchem AG's highly toxic cyanamide and Bayer's propineb, which damage sexual function and fertility.

A total of 230 active ingredients are registered in Kenya, including 51 that are no longer permitted in the EU, such as atrazine (Syngenta), trichlorfon (Bayer) and fipronil (BASF). 70 percent of the rural economy works in the agricultural sector. NGOs warn that farmers are increasingly using dangerous substances to grow food. Despite being banned in the EU, Kenyan imports in 2018 and 2019 included iprodione and acetochlorines from Belgium and 1,3-dichloropropene from Spain. South Africa imported active substances such as imidacloprid, which is hazardous to bees, from Germany and France in 2021 and 2022.


The pesticide companies claim their products are safe and do not endanger humans, insects, or water bodies when handled properly. Proper handling often includes wearing personal protective equipment and adhering to specific application times, spraying distances, and guidelines for co-application with other substances. In reality, the prescribed application often cannot be guaranteed in the Global South, because applicants are inadequately trained or not trained at all, and insufficiently informed about health hazards and distance requirements when applying pesticides. Personal protective equipment is often difficult to obtain, too expensive, or it is simply not reasonable to wear due to high temperatures. Different studies also show that many users aren't able to read the instructions, either because they have a low level of school education or because the instructions are not written in the common languages of the

In the last quarter of 2020, Bayer and Syngenta announced exports of more than 3,800 tonnes of highly hazardous insecticides in third countries like Kenya and Brazil



PLENTY OF MARKETS WITH HIGH MARGINS

The three largest European exporters and the three largest importers of pesticides without EU approval, 2018 in tonnes




country. International organizations such as the FAO and WHO have been pointing out this problem for years.

Human rights experts criticize the practice by EU Member States of exporting EU banned pesticides to the Global South, because it externalizes the health and environmental impacts of these hazardous substances on the most vulnerable. Civil society organizations therefore demand a legal ban of such practices. Pesticides not approved in the EU due to their unacceptable health or environmental effects should no longer be allowed to be sold to countries outside the EU. In 2020 the European Commission's draft chemicals strategy included for the first time a commitment to prevent the export of hazardous chemicals banned in the EU. A first legal draft is to be expected in 2023.

Some European states have already taken national action. In France, a law forbidding the manufacture, storage, and export of EU banned pesticides came into force in January 2022. These substances can no longer be used to maintain green spaces, pathways or forests. Switzerland has banned the export of five particularly toxic pesticides since 2021, with other active ingredients to follow. In Germany, an announcement of putting a legal stop to such exports in the future was confirmed and concretised in September 2022. Importing countries have also taken steps against double standards in pesticide trade: Tunisia, Mexico and the Palestinian National Authority have imposed a ban on imports of pesticides that are forbidden in the exporting or producing country itself. ●

Random samples reveal: As long as it is allowed to export banned pesticides, they will return to Europe – in our fruit and vegetable

If ratified, the EU-Mercosur agreement would reduce over 90 percent of existing tariffs on pesticides and could increase exports of hazardous pesticides from the EU to South America



GLYPHOSATE

A MANUFACTURED CONTROVERSY

Bayer and other companies are fighting for the re-approval of glyphosate in the EU. To do so, they must prove that their pesticide active substance is not carcinogenic. But the studies presented are old – and point to the opposite.

In December 2019, the German pharmaceutical and biotechnological company Bayer submitted an application for re-approval of glyphosate for the European Union (EU) in conjunction with other companies under the name Glyphosate Renewal Group (GRG). Glyphosate is a chemical compound that works as a weed killer. It's the most commonly used herbicide chemical in the world. The approval process is accompanied by a yet unresolved controversy between EU authorities and the World Health Organization's International Agency for Research on Cancer (IARC), which centers on glyphosate's toxicity. In 2015, the IARC had classified the chemical as "probably carcinogenic to humans". The German Federal Institute for Risk Assessment (BfR) and the European Food Safety Authority (EFSA) – both in charge in the EU approval process at the time – came to a different conclusion. As a result of this heated debate, the EU renewed the license for the weed killer for five years, ten years less than the usual authorisation for crop protection chemicals. Glyphosate is currently approved for use as an active ingredient in pesticide products in the EU until the end of 2022.


Bayer's application calling for re-approval is substantiated with hundreds of manufacturer studies and studies from scientific literature but does not contain any new studies refuting the classification of glyphosate as "probably carcinogenic" by IARC. Instead, the Glyphosate Renewal Group relies on twelve cancer studies with mice and rats commissioned by manufacturers, which the agrochemical

company Monsanto – acquired by Bayer in 2018 – had already submitted in the previous approval process.

The IARC evaluated the weight of the evidence and examined four of these twelve cancer studies used by the authorities at the time as evidence of the safety of glyphosate. Basing their judgement exactly on these studies which manufacturers are trying to prove the harmlessness of glyphosate, the WHO cancer researchers concluded "sufficient evidence of carcinogenicity in animal studies". As it later turned out, the BfR had ignored statistically significant increases in tumours in all cancer studies commissioned by manufacturers – according to current rules, two independent studies with positive cancer findings are sufficient to classify a substance as carcinogenic. The BfR justified this failure in an addendum to its assessment report and state that it relied on the statistical evaluations of the manufacturers' study reports. This means that the German authorities did not evaluate the studies' results itself, although its legal mandate is predicated on scientific independence.


Even after being alerted to this fact, the authorities maintained their original conclusion. However, their explanation for considering glyphosate not carcinogenic has changed. The pesticide active ingredient was not responsible for numerous significant tumour findings, they claim, but deficiencies in the conduct of the studies: high dosages, sick laboratory animals, or mere coincidence. It remains questionable how authorities could make an objective assessment of cancer risk based on flawed studies. And secondly, why the manufacturers did not submit new and less flawed cancer studies for the current renewal process.

In the last decade, the U.S. Environmental Protection Agency (EPA) has drastically raised glyphosate tolerances. Civil society organizations state that the EPA is missing key pieces of information including an ecological risk assessment



CONSENSUAL THEFT

Plagiarized and copy pasted content in the chapters on published studies in the 2015 glyphosate report of the German Federal Institute for Risk Assessment (BfR)



But not just the manufacturers' cancer studies have come under criticism. The authorities and IARC also reached different conclusions on the genotoxicity of glyphosate. Based on 53 studies commissioned by manufacturers, the EU authorities in 2015 denied that the herbicide can cause DNA or chromosomal damage. However, similar independent studies from scientific literature – which in their majority support a conclusion of "strong evidence of genotoxicity" according to IARC – had been classified by the EU authorities as "not reliable" and were excluded from the assessment. In September 2017, a plagiarism report revealed that the BfR's declaration in which the regulator had justified the exclusion of these studies was a copy of Monsanto's application for approval. Experts also criticize that national authorities like the BfR just focused on certain aspects like dietary exposures and risks to the general population – leaving risks of occupational exposure out of the picture.

A 2019 ruling by the European Court of Justice requires EU regulators to disclose all manufacturer commissioned studies which had previously been confidential upon request. Two renowned researchers from the Institute of Cancer Research at the Medical University of Vienna in Austria examined the 53 manufacturer commissioned studies mentioned above and evaluated their scientific quality: 34 studies showed substantial deviations from applicable OECD test guidelines and were classified by the two researchers as "not reliable". As for the rest of the 53 studies, 17 were classified as "partly reliable" and only 2 studies as "reliable". However, the applicants resubmitted these studies in the current approval process again as evidence of the lack of genotoxicity of glyphosate.


Notwithstanding all this, in its first draft report of June 2021 the Assessment Group on Glyphosate was proposing to classify glyphosate in the EU as non-carcinogenic and non-toxic again. The group – consisting of EU Member States France, Hungary, the Netherlands and Sweden – is appoint-

Ctrl+C: The German Federal Institute for Risk Assessment has copied entire paragraphs and pages of running text from industry dossiers. A plagiarism report has concluded that the institute even copied the evaluations of independent studies by Monsanto

ed by the European Commission and mandated to ensure that the application meets the formal requirements of the EU legal provision. The EU licence for glyphosate was due to expire in December 2022, but a decision on its future has been pushed back to mid-2023 following an "unprecedented" number of responses to public consultations. ●

MIGHTY INFLUENCER


EU lobbying expenditures of key companies in 2020, in euros



Monsanto's **officially** claimed budget in 2016/17



Monsanto's **actual** lobbying budget in 2016/17



According to Transparency International, the ratio of member of European Parliament to lobbyists is one to fifty. Many of them are sent by agrochemical companies

GENDER

AT THE FOREFRONT OF EXPOSURE

Women working in agriculture often have lower levels of income and lack decision-making power. There is urgent need for gender equality to achieve food security and protection from pesticide exposure.

Women make up 43 percent of the global agricultural labour force, with almost 70 percent of employed women in South Asia and more than 60 percent of employed women in Sub-Saharan Africa working in agriculture. However, women’s participation in agriculture is likely underestimated. Subsistence agriculture, unpaid family work, and seasonal labour, which frequently involve women and girls, often go unaccounted for.

Be it in subsistence farming, informal or formal employment, women are routinely exposed to toxic pesticides. Women carry out a significant part of pesticide application in certain countries and sectors, for example on coffee and fruit farms in South Africa, banana plantations in Costa Rica, or in Malaysia, where there are an estimated 300,000 women sprayers in the plantation sector. A study found that women plantation workers in Indonesia, Malaysia, and the Philippines are frequently exposed to Highly Hazardous Pesticides (HHPs) through mixing, loading, and spraying pesticides. Employers often do not provide Personal Protective Equipment (PPE), so women improvise

by wrapping scarves around their faces or using bra cups as masks or respirators.

Women can also be unknowingly exposed to pesticides through activities like weeding and harvesting which does not require PPE. Women in flower farms in Kenya are more involved in weeding, flower cutting and packaging and showed a higher frequency of poisoning symptoms than men that do the actual spraying.

Recent figures on unintentional acute pesticide poisoning estimate that 385 million or roughly half of the world’s farmers and farmworkers are poisoned each year. However, there is insufficient data to estimate the incidence of poisoning for women because there is a lack of gender-disaggregated data and gender perspective in occupational health research.

Due to traditional gender roles, women are more exposed to pesticides through household chores such as washing spraying equipment or their husbands’ pesticide-soaked clothes, storing pesticides, or disposing pesticide containers. In Vietnam, a study found that more girls reported exposure to pesticides from washing spraying tanks compared to boys.

Pesticides are supposed to prevent crop losses. But a large proportion of losses occur because of inadequate extension, financial support and lack of equipment, especially for women


A MATTER OF SOCIAL JUSTICE

Yield gap between men and women farmers in Uganda




Due to differences in resource use, the worldwide yield gap between men and women farmers averages around 20 to 30 percent. Given equal access to resources, women could achieve the same yield levels. This could reduce the number of undernourished people in the world by 12 to 17 percent and boosting agricultural output in poor countries by 2.5 to 4 percent, according to a NGO study from 2016

Resources for high-yield agriculture: Who benefits?



DUE TO EDUCATION LEVELS, WOMEN ARE AT HIGHER RISK

Gender disparities in pesticide use in Nepal (South Asia), and Ghana (West Africa), in percent



Studies in Bolivia, South Africa, and Tanzania also reveal that lower literacy rates and limited access to training increase women’s vulnerability to pesticides. Women were unable to identify the names of the pesticides they were using, and unable to read or understand safety information on labels.

The impact of pesticides on women and girls differ from the impact on men and boys. Women generally have a higher proportion of body fat, and are thus more likely to store pollutants that can bioaccumulate in fat tissue. Women have a higher level of hormonally sensitive tissues that make them more vulnerable to pesticides, especially those that are hormonally active or known to disrupt the endocrine system. There is an established link between breast cancer and certain pesticides, which act as mammary carcinogens and tumour promoters. Residues of organochlorine pesticides, which degrade slowly and bioaccumulate in the food chain, including banned pesticides such as DDT, have been found in women breast cancer patients. Pesticides are also linked to endometriosis, a painful condition that may cause infertility and can pose a significant risk to women’s reproductive health and their unborn child. Passed on from mother to child through the womb and breastfeeding, pesticides are linked to neonatal deaths, birth defects, and impaired mental development or pervasive developmental problems in children. Studies in the emerging field of epigenetics also show that pesticides exposure may affect gene activity and affect inherited physiological traits.


Women are recognized as playing a key role in transitioning to agroecology – and rural women in the Global South have taken the lead in eliminating pesticides use. Such movements are of benefit not only to farmers, but future generations whose welfare rests upon the health and well-being of women.

Access to land is often denied to women. For many in Latin America, inheritance is the only way to acquire land

More than 80 percent of male cocoa farmers in Ghana posses at least a primary school education certificate, while almost half of female workers at cocoa farms in Ghana have no formal education at all. Studies show how education levels correspond with hazard awareness

UNEQUAL OWNERSHIP AND UNEQUAL OPPORTUNITIES

Form of acquisition of land ownership in Latin America by gender, in percent



DIGITAL TRANSFORMATION

WHO REALLY BENEFITS FROM DIGITALIZATION?

Agricultural robotics, drones, and algorithm-driven technologies for a new way of farming are becoming big business. They are supposed to help farms cut pesticide use, but there are serious doubts whether this will work.

Agriculture faces major challenges. For one thing, it still has to cope with plant diseases, insect pests, and weeds. And secondly, high consumption of pesticides leads to entirely new risks for both humans and nature. Agricultural technology companies are promising to solve these problems with digital technologies known as smart farming or precision farming. According to a survey, 82 percent of farms in Germany already use digital technologies. 45 percent of the farmers surveyed work with GPS-controlled agricultural machinery and 40 percent use agricultural apps for their smartphones or tablets. 32 percent use IT solutions to apply crop protection products or fertilizers to their fields. The networked agriculture market is expected to grow from \$1.8 billion in 2018 to \$4.3 billion by 2023, at an annual growth rate of 19.3 percent during the forecast period. Expectations are high: Progressive digitalization is hoped to enable the world's farms to produce food for nine billion people. Some experts predict digital transformation will raise incomes and protect climate and biodiversity by enabling more precision in pesticide and fertilizer usage – which could lead to lower doses. Digital technologies can also save time which could be used for more labour-intensive methods of pesticide-free cultivation.

One example of the digitalization of agriculture is GPS camera technology. It identifies field areas with weed infestations, so the connected field sprayer opens its nozzles in this section only. Selfpropelled spraying robots use this technology to detect, target, and remove weeds. Drones can


be programmed to spot weed nests from the sky. Algorithms can identify and locate diseased or pest infested plants. According to the manufacturers, all this will soon be part of daily farming business.

In a joint trial project, German companies Südzucker AG and the agricultural technology company Amazone in cooperation with the Danish field robot manufacturer FarmDroid are testing how the use of herbicides and insecticides in sugar beet fields can be reduced. The field robot first sows sugar beet seeds in a precise grid using its GPS system. The robot knows the exact position of the beets and hoes next to and between the rows to remove weeds. In the immediate vicinity of the plant, it is difficult to remove all weeds mechanically without damaging the beet, so the robot sprays agrochemicals right next to the beet, which destroys even the last weeds.


Already today, agricultural machinery can identify how well soils are supplied with nutrients. This information can be fed into cropping plan databases to calculate the necessary amount of fertilizer and pesticides to be applied. Big data corporations are playing a significant role in the development and dissemination of the technology, the processing and the use of the data collected. Google for example works with agencies such as the U.S. National Oceanic and Atmospheric Administration (NOAA). The company wants to use its artificial intelligence programs and the weather agency's vast amounts of data to enable extremely accurate weather forecasts in the future.

Whether the ecological effects of digitalization will be positive or negative depends on many factors. Researchers

The survey in the Russian Astrakhan region shows: Farmers in poorer countries fear being left behind by digital technology – if there is no support



CONCENTRATION STRATEGIES
World's biggest companies in pesticides, seeds and farm machinery, revenues in 2020, in billion euros



see potential to reduce pesticide use. On the other hand, there are also so-called rebound effects, for example increased energy consumption due to new technologies or the expansion of intensive production on land previously used only extensively or not at all, or that is ecologically valuable. There is also a risk that smallholder farmers in lower income countries are excluded from this transformation. They may lack access and knowledge to new technologies. Furthermore, many digital tools are only economical when used at large scale.

This could reinforce monopolization and concentration. One example is the market for agricultural machinery. In 1994, the four largest companies controlled less than one-third of the market – after twenty years of consolidation, they already controlled more than half. Players like John Deere are now staking out their territory through collaborations with agrochemical companies. In the past, the company has already cooperated with pesticide manufacturers such as Syngenta, Dow Agrosiences, BASF and Bayer. Other companies such as CNH Industrial and AGCo have also entered into joint ventures. Venture capital interest in software agricultural technologies is rising as well: From 223 billion US dollars in 2015 to more than 700 billion US dollars in 2017.


Civil society organizations warn of a loss of food sovereignty. New tools and techniques are turning land that is currently managed by smallholder families into agro-industries' profits.

Regulation is needed so that not only corporations benefit from digital farming, but also people and the environment

The market for digital agricultural technology is promising – especially for large corporations that are based in the Global North already dominating other sectors of agriculture

One of the future challenges for policymakers is therefore to prevent the commercialization of climate, nutrition, and crop data and to reinstate farmers sovereignty over their data. Otherwise, there is a risk that digital transformation will contribute to further dependence on unsustainable agriculture. ●

PROFIT FOR WHOM?
Estimated addressable market for precision farming worldwide by 2025, by application (in million US dollars)



EU POLICIES

OBJECTIVES ARE NOT ENOUGH

Even though the EU’s pesticide legislation is the most sophisticated, it has failed to achieve a reduction in use. The EU’s Farm to Fork Strategy seeks to change that. Many questions remain.

Pesticides are high on the agenda at the European level. In its Farm to Fork Strategy from May 2020, the European Commission committed to the objective of reducing the use and risks of synthetic pesticides by 50 percent until 2030, the use of the most hazardous substances by 50 percent, and to introduce a new regulation to reach that goal. The “Save Bees and Farmers” European Citizens’ Initiative, which gathered over 1.2 million signatures, demands an even higher reduction of 80 percent by 2030, a complete phase-out by 2035, and strong support to be given to farmers in their transition towards agroecology.

The current policy to bring down pesticide use, the “Sustainable Use of Pesticides Directive”, was introduced in 2009. The legislation aimed to limit the use of pesticides by promoting alternative practices like Integrated Pest Management (IPM). IPM principles give priority to preventative measures and biocontrol. Biopesticides and as a last step synthetic pesticides are only an option, when all other measures have failed.


However, more than a decade after the adoption of the directive, the EU Court of Auditors (ECA), the Union’s external auditor to assess among other things the effectiveness of EU action, found that only limited progress has been achieved

in measuring and reducing the use and risk of pesticides in the EU. Over the period 2011 to 2018, the sales of pesticides remained stable at around 360 thousand tonnes per year in the EU. The ECA outlined several flaws in the EU framework. For example, there is the missing alignment between the Common Agricultural Policy (CAP) and reduction policy. The CAP determining the funding and priorities of EU agriculture. Another key issue is the lack of appropriate indicators on European level to measure the potential decrease in pesticides. Indicators are mainly based on sales data of pesticides and therefore do not take into account the agriculture area, the volume or the way these substances are used.

From the EU budget, farmers receive financial support based, for the most part, on the number of hectares of the farm. Currently, the EU does not tie the receipt of this EU funding to the respect of IPM principles and other rules laid out in the pesticide directive. This is unlikely to change substantially with the latest attempt to reform the CAP, entering into force in 2023.

Finally, the way Member States handled the implementation of the EU framework into national rules, is another reason for its limited success. Several Member States delayed the transposition into national law and were slow to develop national action plans to implement concrete measures. Civil



At least officially, the EU encouraging with its strategies like “Farm to Fork” natural pest control mechanisms. Integrated pest management is one sort of a sustainable non-chemical methods in agriculture



BANS NOT WITHOUT REASON

Banned Highly Hazardous Pesticides (HHPs) in 2022, in the European Union and selected African States

The most frequently banned pesticides worldwide



society organizations furthermore criticised Member States for not using the legroom they have within the CAP to make the use of IPM more attractive. Member States can use funding from the CAP to offer voluntary schemes that incentivize techniques which lead to a lower use of synthetic pesticides. However, these schemes fail to take a systematic approach needed to reduce pesticides.

According to a joint statement of over 70 civil society organizations, the new EU legislation needs to address all of these concerns to be effective in transforming the food and agriculture system to protect citizens’ health, biodiversity and the climate. They also demand that the regulation entails ambitious and legally binding reduction targets at both EU and national levels, a complete phase-out of the most hazardous pesticides and the use of damaging practices, like aerial spraying or seed coating, as well as a strengthened definition of IPM. Such practices are part of the transformation towards agroecology.

There are also discussions about the role of the EU when it comes to the use of pesticides in other countries. In its Chemicals Strategy for Sustainability from October 2020, the European Commission commits to put an end to pesticides banned in the EU being exported by EU companies to other parts of the world. But this has yet to be translated into actual policies.

The timeline for the reform on the pesticide directive was originally expected at the beginning of 2022 – but was

The global comparison shows: The European Union leads the way in banning very harmful pesticides. But toxic substances are still used in Europe

delayed to summer 2022 due to Russia’s war of aggression against Ukraine. The co-decision between the European Parliament and the Council are expected to go well into 2023 with the new rules likely only being made applicable starting 2024. ●

MONEY, FOR NOT MANY

Value of the pesticide market, in billion euros



The pesticide market has almost doubled in the last 20 years worldwide, with the European Union as one of the biggest consumer and exporting markets

PESTICIDE-FREE REGIONS

GOOD EXAMPLES

All over the world, projects are proving that an ecological future is possible: More and more cities, states, and regions are trying to slash pesticide use; or even ban chemical agents completely from their fields and land.

More than 550 German cities and municipalities have so far decided to manage their urban greenery partially or completely without pesticides. Some municipalities are phasing out a specific group of active ingredients or a specific active ingredient, such as glyphosate. Other municipalities have already completely cut the use of pesticides. One example is Saarbrücken, capital of the German state Saarland: The city has not used pesticides for 25 years. Many cities and regions in the European Union (EU) have also established pesticide-free zones – in Italy, Belgium, the Netherlands, and Luxembourg. However, so far this only affects municipal areas. Many farms in those re-


gions continue to use pesticides. In 2007, Denmark implemented a nationwide ban on the use of pesticides in public areas. Additionally, Danish politicians have been working to reduce the usage of pesticides all over the country. Denmark has slashed nationwide pesticide use by more than 40 percent since 2011. It currently applies an average of 40 percent fewer pesticides than its EU neighbors. Despite these efforts, the country is still far from being completely pesticide-free.

One of the European pioneers in banning pesticides is Luxembourg, where a complete pesticide ban on public land came into force in 2016. Since 2021, the government has also forbidden the use of glyphosate on agricultural land – even though the herbicide is still approved throughout the EU until 2022. The Italian commune of Mals in South Tyrol – the largest apple growing region in Europe – is also particularly committed to living and doing business without harmful pesticides. In a referendum in 2014, the majority of residents decided that their community areas and agricultural land should be pesticide-free. Apart from broad support, the resolution faced a lot of opposition from business, such as large local apple orchards fighting in court to prevent the pesticide ban from being implemented. The administrative court finally overturned the referendum with the argument that the municipality was not the competent authority for this environmental protection issue.

Nevertheless, the civil society effort has received widespread recognition: In 2020, the community was honored with the EuroNatur award for its perseverance in taking action against pesticides.


Not only in Europe, but all over the world a change is taking place. In 2018, Mexico was admonished by the National Human Rights Commission for violating its due diligence obligations by failing to ban Highly Hazardous Pesticides (HHPs). Only two years later, the Mexican agriculture department has proposed rules for phasing out the use of glyphosate by 2024, following pressure from civil society organizations. Until then, a transition period will be established to achieve the total substitution of the herbicide. The competent authorities were urged to develop non-chemical alternatives to current pesticides. Kyrgyzstan even plans to completely phase out pesticide use. Kyrgyzstan’s parliament decided in 2018 that all agriculture should transition to organic production within the next ten years, eliminating the use of synthetic chemical insecticides, herbicides, fungicides, other agricultural chemicals as well as growth regulators. Only biological substances are excluded from the decision. In India, several states have begun to convert their agriculture to or-

Organic farms in Asia, Africa, and Latin America mostly manage small areas. Australia has the largest organically farmed land – more than 35 million hectares



IT IS ABOUT TIME

Share of organic farming on agricultural land in the European Union, by Member State in 2019



ganic farming and ban pesticide use: The small state of Sikkim will be the first region in the world to have a 100 percent organic agriculture. This is a huge paradigm shift in a country that for decades had relied on the heavy use of synthetic fertilizers and pesticides.


Key to the decision in Sikkim were rising cancer rates, polluted rivers, and infertile soils due to pesticide usage. The Sikkim government also attributed its move to the fact that pesticide residues – including many that are banned in other countries – have contaminated staple foods such as rice, vegetables, and fish. The Indian state of Andhra Pradesh – about the size of Austria, Denmark and the Netherlands combined – announced in 2018 that the state’s approximately six million farmers will work without synthetic chemical pesticides by 2024 at the latest. Sri Lanka is following this lead: To achieve the goal of 100 percent organic agriculture, the government had temporarily banned the import of chemical fertilizers and pesticides in April 2021. A few months later after an economic crisis, the government reversed the decision and allowed imports again.

However, the country is sticking to the fight against toxic substances: For years now, the government has been tightening restrictions with the pesticide control law, banning a total of 36 Highly Hazardous Pesticides. For this ef-

IPM is an approach to suppress pest populations. It uses biological and ecological knowledge to avoid pesticides – their use is a last resort

Member States still have until 2030 to increase organically farmed land to a quarter of total agricultural land, as stipulated in the EU’s Farm to Fork Strategy

fort, the country received the Special Future Policy Award in 2021, dedicated to the most effective policy solutions that protect people and the environment from hazardous chemicals. ●



AUTHORS AND SOURCES FOR DATA AND GRAPHICS

All online links were last checked in November 2022.
See page 2 for the websites where you can download a clickable PDF of this atlas. Lengthy links have been shortened using the bitly web address conversion service.

08–09 BRIEF LESSONS

Nyamu, D. et al., 2012, Trends of Acute Poisoning Cases Occurring at the Kenyatta National Hospital, <https://bit.ly/3ChYsdX>. RTFI & EcoTrac, 2022. Pesticide Use in Kenya: A Toxic Business. Heinrich Böll Stiftung Nairobi Office (unpublished). Kenya Organic Agriculture Network, 2020, Pesticide Use in Kirinyaga and Murang’a Counties, <https://bit.ly/3dyCXfv>. Route to Food Initiative, 2019, Pesticides in Kenya: Why Our Health, Environment and Food Security are at Stake, <https://bit.ly/3y0HAWQ>.

10–11 PESTICIDES AND AGRICULTURE DANGEROUS SUBSTANCES by Lisa Tostado

p.12: Company reports, quarterly statements. Jennifer Clapp, The problem with growing corporate concentration and power in the global food system, 2021, <https://go.nature.com/3xTA9iR>. – **p.13 top:** Faostat, <https://bit.ly/3DohlJQ>. – **p.13 bottom:** Yijia Li, Ruiqing Miao, Madhu Khanna, Neonicotinoids and decline in bird biodiversity in the United States, 2020, <https://go.nature.com/3Epuof7>.

12–13 CORPORATIONS BIG PROFITS WITH TOXIC TRADE by Carla Hoinkes

p.14: Public Eye, <https://bit.ly/3GrePUS>. – **p.15 top:** Public Eye, <https://bit.ly/3G8awhc>. – **p.15 bottom:** Faostat, <https://bit.ly/3DohlJQ>. PAN Germany, Giftige Exporte. Die Ausfuhr hochgefährlicher Pestizide von Deutschland in die Welt, 2019, <https://bit.ly/3rFELli>.

14–15 PESTICIDE USE IN AFRICA INCREASINGLY VULNERABLE by Cecilia Moraa Onyango

p.16: FAO Statistics and Database of the Food and Agriculture Organization of the United Nations (FAOSTAT), accessed 21st September 2022, <https://bit.ly/2PknqxO>. – **p.17:** Tolera, S., 2020, Systematic review on adverse effect of pesticide on top ten importers of African countries, <https://bit.ly/3Ryy4Ry>. – Jepson, P.C., et al., 2020, Selection of pesticides to reduce human and environmental health risks: A global guideline and minimum pesticides list, <https://bit.ly/3RB7VBH>. – PAN International, 2022, List of Highly Hazardous Pesticides, <https://bit.ly/3SIVwwM>.

16–17 NIGERIA TIME FOR A DETOX IN AGRICULTURE by Silke Bollmohr, Ikenna-Donald Ofoegbu and Jochen Luckscheiter

p.16: Silke Bollmohr, Time for a Detox in Agriculture – Challenges of Pesticide Use and Regulation in Nigeria and Possible Solutions, 2020, <https://bit.ly/3g6kwQp>. – **p.17:** The Vanguard, EU bans Nigeria from exporting beans, melon seeds, dried fish, meat and others, 10.08.2015, <https://bit.ly/3CViT0X>. – Deutsche Welle, Nigeria: Pesticide caused death of nearly 300 villagers, 09.10.2021, <https://bit.ly/3EzfdTH>.

18–19 APPROVAL PROCEDURES UNDERESTIMATED RISKS by Carsten Brühl and Johann Zaller

p.18: European Commission, Guidelines on Active Substances and Plant Protection Products, <https://bit.ly/3GbKKsk> – **p.19:** PA International, Pesticide Use In The EU – Presence Of Candidates For Substitution And Low Risk Active Substances, 2021, unpublished. IBMA Market Survey 2021, <https://bit.ly/3or49zD>.

20–21 HEALTH SEVERE CONSEQUENCES by Wolfgang Bödeker

p.20: Leonardo Trasande et al., Estimating Burden and Disease Costs of Exposure to Endocrine-Disrupting Chemicals in the European Union, 2015, <https://bit.ly/31DeGPv>. – **p.21:** Wolfgang Boedeker et al., The global distribution of acute unintentional pesticide poisoning: estimations based on a systematic review, 2020, <https://bit.ly/3r3Tj41>.

22–23 SOILS IMPACT ON THE INVISIBLE ECOSYSTEM by Johann Zaller

p.22: Vera Silva et al., Pesticide residues in European agricultural soils – A hidden reality unfolded, 2019, <https://bit.ly/3QRdYTm>. – **p.23 top:** Vera Silva et al., Pesticide residues in European agricultural soils – A hidden reality unfolded, 2019, <https://bit.ly/3QRdYTm>. – **p.23 bottom:** Judith Riedo et al., Widespread Occurrence of Pesticides in Organically Managed Agricultural Soils – the Ghost of a Conventional Agricultural Past?, 2021, <https://bit.ly/3ufnsze>.

24–25 RESIDUES IN KENYA TOXIC SIDE DISH by Silke Bollmohr, Susan Haffman and Catherine Kunyanga

The contents of this article have been modified from the original version of the publication. **p.24:** FSA, The 2018 European Union report on pesticide residues in food, <https://bit.ly/3bui4kj>. Kenya Plant Health Inspectorate Service, 2018, Annual Report. Kenya Organic Agriculture

Network, 2020, Pesticide Use in Kirinyaga and Murang’a Counties, <https://bit.ly/3dyCXfv>. – **p.25 top:** Public Eye, <https://bit.ly/3rD1l4j>. – **p.25 bottom:** Silke Bollmohr, 2021, Time for a “Detox” in Agriculture Pesticide Use in Four States in Nigeria published by the Heinrich Böll Foundation Abuja Office, <https://bit.ly/3SuSV9S>.

26–27 BIODIVERSITY EXTINCTION IN FULL SWING by Katrin Wenz

p.26: Tari Gunstone et al., Pesticides and Soil Invertebrates: A Hazard Assessment, 2021, <https://bit.ly/3GhG3NA>. – **p.27 top:** Caspar A. Hallmann et al., More than 75 percent decline over 27 years in total flying insect biomass in protected areas, 2017, <https://bit.ly/3IEvRXP>. Jörg Hoffmann, IPBES, Global Assessment Report on Biodiversity and Ecosystem Services, 2019, <https://bit.ly/3bwEi4Y>. Tim Wahrenberg, Effects of cultivation practice on floristic and flowering diversity of spontaneously growing plant species on arable fields, 2021, <https://bit.ly/3EGKKQR>. – **p.27 bottom:** European Environment Agency, Conservation status of species under the EU Habitats Directive, <https://bit.ly/3OZwg3p>.

28–29 BENEFICIAL INSECTS SMALL CREATURES WITH BIG IMPACTS by Sevgan Subramanian, Saliou Niassy, Samira Mohamed, Chrysantus Tanga and Sunday Ekesi

p.28: Daphne, M., Pollinators in Africa: Understanding is the first step to protecting, 2011, <https://bit.ly/3rBpneU>. – Macharia, I., Löhr, B., & De Groote, H., Assessing the potential impact of biological control of *Plutella xylostella* (diamondback moth) in cabbage production in Kenya, 2005, <https://bit.ly/3CDqWiP>. – Mohamed, S. A., et al., A deadly encounter: Alien invasive *Spodoptera frugiperda* in Africa and indigenous natural enemy, *Cotesia icipe* (Hymenoptera, *Braconidae*), 2021, <https://bit.ly/3ElHpce>. – Niassy, S., et al., Insight on fruit fly IPM technology uptake and barriers to scaling in Africa, 2022, <https://bit.ly/3RGcLh7>. – Sagwe, R. N., et al., Pollinator supplementation mitigates pollination deficits in small-holder avocado (*Persea americana* Mill.) production systems in Kenya, 2021, <https://bit.ly/3M8ZZ9N>. – **p.29 top:** Tanga, C. M., et al., Edible insect farming as an emerging and profitable enterprise in East Africa, 2021, <https://bit.ly/3SEJHic>. – Zeddies, J., et al., Economics of biological control of cassava mealybug in Africa, 2001, <https://bit.ly/3C5HGgQ>. – **p.29 bottom:** Naturkapital Deutschland – Teeb De, Ökosystemleistungen in ländlichen Räumen, 2016, <https://bit.ly/3IKdMY7>, S. 103, Abb. 5.12.

30–31 WATER GO WITH THE FLOW by Silke Bollmohr, based on an article by Falk Hilliges, Kristina Hitzfeld, Jan Koschorreck and Alexandra Müller

p.30: Jorge Casado et al., Screening of pesticides and veterinary drugs in small streams in the European Union by liquid chromatography high resolution mass spectrometry, 2019, <https://bit.ly/3rxacDe>. European Commission,

Manure and soil biodiversity, <https://bit.ly/3njj4u8>. European Environment Agency, <https://bit.ly/3QR2Gi0>. <https://bit.ly/3A65UbM>. <https://bit.ly/3ngy0cu>. – **p.31 top:** Bund/Länder-Arbeitsgemeinschaft Wasser, Bericht zur Grundwasserbeschaffenheit, 2019, <https://bit.ly/3IAUTH3>. European Environment Agency, <https://bit.ly/3OG07jZ>. – **p.31 bottom:** Life Apex, <https://bit.ly/3DpslX9>.

32–33 LONG-RANGE TRANSPORT GONE WITH THE WIND by Johanna Bär, Johannes Heimrath and Anna Satzger

p.32: Caroline Linhart et al., Year-round pesticide contamination of public sites near intensively managed agricultural areas in South Tyrol, 2021, <https://bit.ly/30WhIxY>. – **p.33 top:** Umweltinstitut München, Pestizid-Belastung der Luft, 2020, <https://bit.ly/3nLnafN>. – **p.33 bottom:** Rapunzel Naturkost, <https://bit.ly/3r5LPhe>. Umweltinstitut München, <https://bit.ly/3l3yUs4>.

34–35 INSECT DECLINE AN ECOLOGICAL ARMAGEDDON by Dave Goulson

p.34: D. Susan Willis Chan & Nigel E. Raine, Population decline in a ground-nesting solitary squash bee (*Eucera pruinosa*) following exposure to a neonicotinoid insecticide treated crop (*Cucurbita pepo*), 2021, <https://go.nature.com/3FkmsfC>. Kiah Tasman, Sean A. Rands, James J. L. Hodge, The Neonicotinoid Insecticide Imidacloprid Disrupts Bumblebee Foraging Rhythms and Sleep, 2020, <https://bit.ly/3e9EGVi>. – **p.35 top:** IPBES, The assessment report on pollinators, pollination and food production, 2017, <https://bit.ly/3DlpYoo>. Fiona H. M. Tang et al., Risk of pesticide pollution at the global scale, 2021, <https://go.nature.com/2ZWSoXN>. Edward A. D. Mitchell et al., A worldwide survey of neonicotinoids in honey, 2017, <https://bit.ly/2ZXoXVX>. – **p.35 bottom:** Francisco Sánchez-Bayo, Kris A.G. Wyckhuys, Worldwide decline of the entomofauna: A review of its drivers, 2019, <https://bit.ly/3dlbpXg>. IPBES, The assessment report on pollinators, pollination and food production, 2017, <https://bit.ly/3DlpYoo>.

36–37 HIGHLY HAZARDOUS PESTICIDES A GLOBAL HUMAN RIGHTS CONCERN by Silke Bollmohr, Susan Haffman

p.36: EcoTrac, Pesticides in Kenya: Where are we?, <https://bit.ly/3lGA65b>. – **p.37 top:** PAN International, List of Highly Hazardous Pesticides, 2022. – **p.37 bottom:** PAN Germany, Giftige Exporte, 2019, <https://bit.ly/3AILjQT>.

38–39 GENETIC ENGINEERING MORE MODIFIED CROPS, MORE PESTICIDES by Martha Mertens

p.40: IHS Markit, Analysis of sales and profitability within the seed sector, 2019, <https://bit.ly/3pvOsQ8>. Jennifer Clapp, The problem with growing corporate concentration and power in the global food system, 2021,

<https://go.nature.com/3xTA9iR>. – **p.41:** International Service for the Acquisition of Agribiotech Applications, <https://bit.ly/3do7aKC>. Faostat, <https://bit.ly/3DohlJQ>. The International Survey of Herbicide Resistant Weeds, <https://bit.ly/3Igjwhr>.

**40–41 SMALLHOLDER FARMERS
NEW MARKETS, LESS REGULATIONS**
by Layla Liebetrau

The contents of this article have been modified from the original version of the publication.
p.42: Agrochemical Association of Kenya, Activity Report April to June 2022, <https://bit.ly/3UeWsdD>. – **p.43 top:** David Guereña, Supporting smallholders in maintaining soil health: key challenges and strategies, 2018, <https://bit.ly/3duLaxJ>. Philipp Staudacher et al., Comparative Analysis of Pesticide Use Determinants Among Smallholder Farmers From Costa Rica and Uganda, 2020, <https://bit.ly/3rKqVob>. European Union, The use of pesticides in developing countries and their impact on health and the right to food, 2021, <https://bit.ly/3y11wYo>. Vu Ngoc Huyen et al., Effects of pesticides on farmers’ health in Tu Ky district, Hai Duong province, Vietnam, 2020, <https://bit.ly/3duEWOH>. Ulrike Bickel, Uso de plaguicidas por productores familiares en Bolivia, 2018, <https://bit.ly/3EzAmu9>. EuRh, Nachhaltige Verwendung von Pflanzenschutzmitteln: begrenzter Fortschritt bei der Messung und Verringerung von Risiken, 2020, <https://bit.ly/3y2jidL>. – **p.43 bottom:** Philipp Staudacher et al., What agro-input dealers know, sell and say to smallholder farmers about pesticides: a mystery shopping and KAP analysis in Uganda, 2021, <https://bit.ly/3xU3pG4>.

**42–43 IMPORTS AND EXPORTS
BANNED BUT SOLD ANYWAY**
by Silke Bollmohr and Susan Haffman
p.44: IPBES, The assessment report on pollinators, pollination and food production, 2017, <https://bit.ly/3DlpYoo>. – **p.45 top:** Public Eye, <https://bit.ly/332kKBw>. – **p.45 bottom:** Greenpeace Austria, Pestizide in brasilianischem Obst, 2020, <https://bit.ly/3bwKvOE>. Greenpeace Germany, Pestizide aus Deutschland in brasilianischem Obst, 2021, <https://bit.ly/3lO5w9w>. Public Eye, <https://bit.ly/3dTlyhG>.

**44–45 GLYPHOSATE
THE MANUFACTURED CONTROVERSY**
by Helmut Butscher-Schaden
p.46: As You Sow, Roundup revealed: Glyphosate in our Food System, 2017, <https://bit.ly/3u6UCjt>. Global Industry Analysts, Inc, <https://bwnews.pr/3oCP1iI> – **p.47 top:** Stefan Weber, Helmut Burtischer-Schaden, Detailed Expert Report on Plagiarism and superordinated Copy Paste in the Renewal Assessment Report (RAR) on Glyphosate, 2019, <https://bit.ly/331J8TR>. Armen Nersesyan, Siegfried Knasmueller, Evaluation of the scientific quality of studies concerning genotoxic properties of glyphosate, 2021, <https://bit.ly/3IlqK8A>. – **p.47 bottom:** Corporate Europe Conservatory, <https://bit.ly/3onRTzA>. LobbyFacts, <https://bit.ly/33drZGG>.

**46–47 GENDER
AT THE FOREFRONT OF EXPOSURE**
by Ilang-Ilang Quijano
p.48: Farming First, <https://bit.ly/2O40bfq>. – **p.49 top:** Kishor Atreya, Pesticide use knowledge and practices: A gender differences in Nepal, 2007, <https://bit.ly/3buPMGr>. Yaw Osei-Owusu, Raymond Owusu-Achiaw, Assessment on the Gender Dynamics of Highly Hazardous Pesticides (HHPs) within Cocoa Production Landscape in Ghana, <https://bit.ly/3fLg0Xm>. – **p.49 bottom:** Carmen Diana Deere and Magdalena León, The Gender Asset Gap. Land in Latin America. World Development 31, 2003, <http://bit.ly/1GXyXuH>. FAO, The State of Food and Agriculture: Women in Agriculture, 2011, <https://bit.ly/3ykEJs6>.

**48–49 DIGITAL TRANSFORMATION
WHO REALLY BENEFITS FROM DIGITAL
TRANSFORMATION?**
by Heike Holdinghausen
p.50: Aleksandr Koshkarov, Tatiana Koshkarova, Data-driven approach in digital agriculture: survey of farmers, 2019, <https://bit.ly/3nk7qiC>. – **p.51 top:** Company reports, quarterly statements. Jennifer Clapp, The problem with growing corporate concentration and power in the global food system, 2021, <https://go.nature.com/3xTA9iR>. Pat Mooney, Blocking the Chain, 2018, <https://bit.ly/3rMVU2T>. Market Data Forecast, <https://bit.ly/31E82rT>. **p.51 bottom:** BIS Research, Precision Agriculture Market, <https://bit.ly/3QP9bSn>.

**50–51 EU POLICIES
OBJECTIVES ARE NOT ENOUGH**
by Clara Bourgin and Andre Prescher
p.52: Jacob R. Pecenka et al., PM reduces insecticide applications by 95% while maintaining or enhancing crop yields through wild pollinator conservation, 2021, <https://bit.ly/3HUesV1>. – **p.53 top:** PAN International, List of Highly Hazardous Pesticides, 2022. – **p.53 bottom:** Bureau for the Appraisal of Social Impacts for Citizen information, A Model that’s costing us dearly, 2021, <https://bit.ly/3QHUpgt>.

**52–53 PESTICIDE-FREE REGIONS
GOOD EXAMPLES**
by Ulrike Bickel
p.54: FiBL & IFOAM, The World of organic agriculture, 2019, <https://bit.ly/34rsnCj>. – **p.55 top:** Eurostat, Für ökologische Landwirtschaft genutzte Fläche, <https://bit.ly/3GjQa4O>. Thünen Report, Leistungen des ökologischen Landbaus für Umwelt und Gesellschaft, 2019, <http://bit.ly/35e6zW3>. **p.55 bottom:** European Commission, <https://bit.ly/3xUyPxe>.

PUBLISHED IN ENGLISH





HEINRICH-BÖLL-STIFTUNG
Our objectives: Fostering democracy and upholding human rights, taking action to prevent the destruction of the global ecosystem, advancing equality between women and men, securing peace through conflict prevention in crisis zones, and defending the freedom of individuals against excessive state and economic power. We are closely tied to the German Green Party (Alliance 90/The Greens), and maintain currently 32 international offices in approximately 60 countries.
Heinrich-Böll-Stiftung,
Schumannstr. 8, 10117 Berlin, Germany, <https://boell.de>

**BUND FÜR UMWELT UND NATURSCHUTZ
DEUTSCHLAND**
We view ourselves as a driving force for ecological renewal, social justice and sustainable development. With more than 660,000 members and supporters, BUND is one of the largest environmental organizations in Germany. We are a member of the Friends of the Earth International (FoEI) network with partner organizations in 72 countries.
Bund für Umwelt und Naturschutz Deutschland (BUND),
Kaiserin-Augusta-Allee 5, 10553 Berlin, Germany, <https://bund.net>

FRIENDS OF THE EARTH EUROPE
We are the largest grassroots environmental network in Europe and campaign on today’s urgent environmental and social issues. We challenge the current model of economic and corporate globalization, and promote solutions that will help to create environmentally sustainable and socially just societies. We advocate for an ecological and fair agriculture that protects natural resources, supports small scale family farms, and halts exploitation of developing countries.
Friends of the Earth Europe,
Rue d’Edimbourg 26, 1050 Brussels, Belgium, <https://foeeurope.org>

**PESTICIDE ACTION NETWORK
EUROPE**
PAN Europe is a network of NGOs working to reduce the use of hazardous pesticides and have them replaced with ecologically sound alternatives. We work to safe sustainable pest control methods. Our network brings together over 45 consumer, public health and environmental organizations and women’s groups from across Europe.
Pesticide Action Network Europe,
Rue de la Pacification 67, 1000 Brussels, <https://pan-europe.info>



Pesticide imports into African countries are increasing. However, there is not enough information about how they are used and the impacts they are having on human health and the environment.

from: **PESTICIDE USE IN AFRICA**, page 14

Surveys have shown that 80 percent of the pesticides used most frequently by small-scale farmers in Nigeria are highly hazardous pesticides.

from: **TIME FOR A DETOX IN AGRICULTURE**, page 16

Beneficial insects do not only protect crops from pests, but they also aid in pollination. Several crops grown in Africa depend on insect pollinators to improve crop yields and quality.

from: **SMALL CREATURES WITH BIG IMPACTS**, page 28